

RESEARCH

Open Access

Genetic basis of pyrethroid resistance in a population of *Anopheles arabiensis*, the primary malaria vector in Lower Moshi, north-eastern Tanzania

Johnson Matowo^{1,4*}, Christopher M Jones², Bilali Kabula^{1,5}, Hilary Ranson², Keith Steen², Franklin Mosha^{1,4}, Mark Rowland^{3,4} and David Weetman²

Abstract

Background: Pyrethroid resistance has been slower to emerge in *Anopheles arabiensis* than in *An. gambiae s.s* and *An. funestus* and, consequently, studies are only just beginning to unravel the genes involved. Permethrin resistance in *An. arabiensis* in Lower Moshi, Tanzania has been linked to elevated levels of both P450 monooxygenases and β -esterases. We have conducted a gene expression study to identify specific genes linked with metabolic resistance in the Lower Moshi *An. arabiensis* population.

Methods: Microarray experiments employing an *An. gambiae* whole genome expression chip were performed on *An. arabiensis*, using interwoven loop designs. Permethrin-exposed survivors were compared to three separate unexposed mosquitoes from the same or a nearby population. A subsection of detoxification genes were chosen for subsequent quantitative real-time PCR (qRT-PCR).

Results: Microarray analysis revealed significant over expression of 87 probes and under expression of 85 probes (in pairwise comparisons between permethrin survivors and unexposed sympatric and allopatric samples from Dar es Salaam (controls). For qRT-PCR we targeted over expressed ABC transporter genes (*ABC '2060'*), a glutathione-S-transferase, P450s and esterases. Design of efficient, specific primers was successful for *ABC '2060'* and two P450s (*CYP6P3*, *CYP6M2*). For the *CYP4G16* gene, we used the primers that were previously used in a microarray study of *An. arabiensis* from Zanzibar islands. Over expression of *CYP4G16* and *ABC '2060'* was detected though with contrasting patterns in pairwise comparisons between survivors and controls. *CYP4G16* was only up regulated in survivors, whereas *ABC '2060'* was similar in survivors and controls but over expressed in Lower Moshi samples compared to the Dar es Salaam samples. Increased transcription of *CYP4G16* and *ABC '2060'* are linked directly and indirectly respectively, with permethrin resistance in Lower Moshi *An. arabiensis*.

Conclusions: Increased transcription of a P450 (*CYP4G16*) and an ABC transporter (*ABC 2060*) are linked directly and indirectly respectively, with permethrin resistance in Lower Moshi *An. arabiensis*. Our study provides replication of *CYP4G16* as a candidate gene for pyrethroid resistance in *An. arabiensis*, although its role may not be in detoxification, and requires further investigation.

Keywords: *Anopheles arabiensis*, Genes, Microarrays, Permethrin, Resistance, Transcription

* Correspondence: johntowo@yahoo.com

¹Kilimanjaro Christian Medical University College (KCMUCo), Moshi, Tanzania

⁴Pan-African Malaria Vector Control Consortium (PAMVERC), Moshi, Tanzania

Full list of author information is available at the end of the article

Background

Vector control programmes employing ITNs and IRS are a mainstay of malaria control [1,2]. Continuous usage of insecticides under such malaria control programmes and/or agricultural application of insecticides has resulted in the development of resistance in major malaria vector species [3-7]. Anopheline mosquito populations in many parts of Africa have developed resistance to pyrethroids [8], which are widely used for indoor residual spraying (IRS) and the only insecticide class available for insecticide treated nets (ITNs). Pyrethroids have been shown to pose very low health risks to humans and other mammals, but are toxic to insects and knock them down (kill them), even at very low doses [9].

Insecticide resistance is considered a major threat to the continued success of IRS and ITNs, with reduction in efficacy of pyrethroids being a particular concern [8,10]. The various mechanisms by which mosquitoes are thought to develop resistance to insecticides include metabolic resistance, target-site resistance, reduced penetration and behavioural avoidance. However, metabolic resistance and target-site resistance are major mechanisms assumed to be responsible for insecticide resistance [11] and are known to contribute to pyrethroid resistance in malaria vectors. While target-site resistance occurs when the site of action of insecticide is altered such that the insecticide no longer binds effectively, metabolic resistance involves over-expression of enzymes capable of detoxifying insecticides or amino acid substitutions within these enzymes, which alter the affinity of the enzyme for the insecticide [12]. The most common target site resistance mechanism constitutes two point mutations at amino acid position 1014 of the voltage-gated sodium channel gene, resulting in either a leucine-phenylalanine (L1014F) [13], or a leucine-serine (L1014S) substitution [14].

Detoxification enzymes include glutathione-S-transferases (GSTs), esterases and P450 monooxygenases. GST-based DDT resistance by breaking down DDT to non-toxic products is common in a number of anopheline species including *An. gambiae* [15-17]. It has been suggested that GSTs may play a role in pyrethroid resistance by detoxifying lipid peroxidation products induced by pyrethroids and/or by protecting from insecticide exposure induced oxidative stress [18]. GSTs might also confer a secondary role in pyrethroid resistance by sequestering the insecticide, hence reducing the total *in vivo* concentration of insecticide [19]. More recently, the study of Riveron *et al.*, 2014 [20] revealed a more direct role played by GSTs in conferring insecticide resistance in malaria vectors.

Elevated levels or activity of esterase enzymes which hydrolyze ester bonds or sequester insecticides is one of the most common metabolic resistance mechanisms in organophosphate (OPs) resistant mosquito species.

Malathion (OP) resistance in *Anopheles culicifacies* and *Anopheles stephensi* has been associated with an altered form of esterase that specifically metabolizes the molecule at a much faster rate than that in susceptible strains [21,22].

Cytochrome P450-dependent monooxygenases are an important and diverse family of enzymes involved in the metabolism of numerous endogenous and exogenous compounds. Cytochrome P450 belongs to six families and increased transcription of genes belonging to the CYP4, CYP6, and CYP9 has been observed in various insecticide-resistant species [23]. Microarray-based approaches have identified three candidate P450 genes, CYP6M2, CYP6P3 and CYP6Z2 that were found to be repeatedly over-produced in pyrethroid resistant populations of *An. gambiae* [24-26]. All of these genes encode for enzymes that are able to bind to type I and type II pyrethroids but only CYP6P3 and CYP6M2 were shown to metabolize the insecticides [27,28]. More recently, some studies demonstrated that CYP6M2 is also capable of metabolizing the organochlorine insecticide DDT in *An. gambiae*, hence demonstrating the first evidence for a metabolic cross-resistance in malaria vectors [29].

Both metabolic resistance and target-site resistance have been documented in *An. arabiensis*, one of the dominant vector species of malaria in sub-Saharan Africa [30,31] and in some areas such as the Great Rift Valley in East Africa, *An. arabiensis* is the predominant malaria vector species [32]. To date, the *kdr* mutations have been reported in several countries including Uganda [33], Sudan [34,35], Cameroon [36], and Tanzania [37-39]. In addition to target site resistance, metabolic resistance has also been documented in different countries [40-42].

The study of dynamics of insecticide resistance in Lower Moshi *An. arabiensis* from 2009 to 2013 [39] has shown that the population has developed resistance to all pyrethroids tested (permethrin, deltamethrin and lambda-cyhalothrin) with the presence of L1014F mutation at very low frequency. Tanzania has been scaling-up the use of pyrethroid-based LLINs which reached its universal coverage under Universal Coverage Campaign [43]. Therefore, higher pyrethroid resistance is expected following higher selection pressure on this Lower Moshi *An. arabiensis* population and it is likely that pyrethroid resistance will spread to other areas in the country. The previous study by Matowo *et al.*, 2010 [44] had revealed significantly higher levels of oxidase and β -esterase enzymes in wild Lower Moshi *An. arabiensis* than in a laboratory reference strain. However, the genes responsible for elevated levels of these detoxification enzymes remained unknown, although over-expression of a few detoxification genes has been documented in *An. arabiensis* populations in Hai district, adjacent to Lower Moshi. [45]. Therefore, the aim of this study was to explore and identify specific

genes involved in metabolic resistance to permethrin in Lower Moshi populations of *An. arabiensis* from Lower Moshi.

Methods

Mosquito sample collection and identification

Collections were conducted in the rainy season between May and June 2011 in Mabogini village (37°21' E, 3°24' S), Mabogini is an agricultural area of about 15 km south of Moshi town, in which rice is cultivated in two growing seasons, the main season in which rice is sown is mid-June with irrigation from June to October, and a second season of sporadic cultivation from September to February. The rice paddies thus provide breeding sites for *Anopheles arabiensis*, the predominant malaria vector in the area, throughout much of the year.

The larvae were collected from two specific localities, Mbugani (37°22' E, 3°25' S) and Harusini (37°21' E, 3°23' S) in Mabogini village which are 4 km. apart. Additional larval collections were made from Mbugani in August 2012.

Anopheline larvae were also collected from temporary breeding habitats in Ilala (Gerezani) (39°16' E, 6°49' S) and Kinondoni (Kawe) (39°13' E, 6°43' S) in Dar es Salaam about 700 km from Moshi, in August and October 2011.

The field-collected anopheline larvae were transferred into an insectary and reared to adults. Emergent adults were split by sex before mating could occur and fed with 10% sugar solution.

Virgin females were preferred following the previous whole-body microarray experiments [46] that detected large transcriptional changes in mated females where the number of genes differentially expressed increased with time post mating.

At 3 days-old, adult females from Mbugani were exposed to 0.75% permethrin for one hour using a standard WHO tube bioassay procedure [47] with mortality scored 24 hours after the exposure period. Adult mosquitoes from Harusini, Moshi and Dar es Salaam were not exposed to insecticide. At their fourth day post emergence, all unexposed (control) females and the survivors (resistant mosquitoes) were killed in 75% ethanol. A hind leg was removed from each mosquito and stored over silica gel in an individually-labelled tube before placing the remainder of the specimens in RNA later in an Eppendorf tube. DNA was extracted from the legs that were removed from each specimen for species identification.

Species identification focused on the most prevalent local vectors of *An. gambiae sensu lato*; *An. arabiensis* and *An. gambiae s.s.* Mosquitoes were identified to species level using standard PCR method [48]. Only those identified as *An. arabiensis* were further analysed in this study.

Whole genome microarrays

Experimental design

The study was designed to comprise of two microarray experiments. The first involved three groups of samples collected in 2011 from Mbugani (insecticide exposed, selecting for the 24% and 54% of the most resistant females [39] and unexposed) and Harusini (unexposed). Although initially suspected that Harusini might display a lower resistance level than Mbugani, its close proximity made predictions ambiguous and only results from hybridizations of Mbugani permethrin-selected against Mbugani unexposed and Mbugani permethrin-selected against Harusini unexposed were used. In the second experiment, to investigate temporal repeatability of results we repeated the experiment from Mbugani (permethrin-selected against unexposed) using a collection from 2012. In this second experiment a pyrethroid-susceptible sample from Kinondoni and Ilala in Dar-es-Salaam [40] was also included, primarily to permit an allopatric comparison with the population from Mbugani (insecticide exposed that selected for the 54% of the most resistant females [39]). However, we found that comparisons involving the Dar-es-Salaam sample yielded very little additional resolution for the analysis, which coupled with suspected technical errors in some of the hybridizations led to their exclusion.

2011: Mbugani permethrin selected vs Mbugani unexposed

2011: Mbugani permethrin-selected vs. Harusini unexposed


2012: Mbugani permethrin-selected vs. Mbugani unexposed

Comparisons in experiment 1 consisted of three independent biological replicates while those of experiment 2 consisted of four independent biological replicates (Figure 1); and two technical repeats of dye swaps to control for dye bias. Hybridization and analysis followed an interwoven loop design.

To be judged significant a probe was required to (a) exhibit a false discovery rate-corrected probability of $q < 0.05$ in all three comparisons, and (b) exhibit a consistent directionality in expression, i.e. either Mbugani permethrin-selected groups > unexposed groups or Mbugani permethrin-selected groups < unexposed groups in all three comparisons. Owing to the use of within population (i) selected vs. (ii) unexposed comparisons, the magnitude of differential expression is expected to be relatively low, because (i) is a subset of (ii) [20] and therefore no fold-change threshold was applied as a significance criterion.

RNA extraction and labelling

Both four day old unfed unmated female mosquitoes that survived one hour exposure to 0.75% permethrin,



held for 24 hours post-exposure and unexposed mosquitoes were killed using 70% absolute ethanol and stored in RNAlater® at -20°C. Total RNA was extracted from batches of ten mosquitoes using the RNAqueous^R-4PCR Kit (Ambion) according to manufacturer's instructions and treated with DNase I (Qiagen). The quality and quantity of the RNA was checked using a Nanodrop spectrophotometer (Nanodrop Technologies, Wilmington, DE, USA) and Bioanalyser (Agilent Technologies, Stratagene, USA) respectively. Extracted mRNA was dye-labelled using the Low Input Quick Amp Labelling Kit (Agilent Technologies, Stratagene, USA) according to manufacturer's instructions. Labelling was repeated for samples that had a low cDNA yield (<15 ng/μl) and poor dye incorporation (<6.0 pmol/μl of each dye).

The labelled samples were hybridized using a Gene Expression Hybridization Kit (Agilent Technologies). Washing, scanning and feature extraction were performed according to the manufacturer's recommendations.

Microarray analysis

Analysis of the microarray data was performed on corrected spot intensities using the LIMMA 2.4.1 software package [49] for R 2.3.0. Normalization of signal intensities was followed by ANOVA F-test using the

MAANOVA package in R [50]. Microarray data are available in the ArrayExpress database (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-2513 (SURV vs UNEXPOSED loop design) and E-MTAB-2514 (SURV vs DAR loop design).

Quantitative real time PCR (qPCR)

Technical repeatability of microarray results for a subset of candidate genes was assessed using Quantitative PCR using the cDNA synthesised from the same RNA samples used in the microarray experiments. The samples were quantified using a Nanodrop spectrophotometer (Nanodrop Technologies, Wilmington, DE, USA). cDNA was synthesised from ~0.5-1 μg of RNA using oligo(dT) 20 (50 μM) and SuperScript III (200U) (Invitrogen) and purified through a DNA-binding column (Qiagen). The quality and quantity of cDNA was measured using the Nanodrop spectrophotometer (Nanodrop Technologies, Wilmington, DE, USA) and the cDNA samples were stored at -70°C until further use.

Primers for qPCR were designed using NCBI primer BLAST (<http://www.ncbi.nlm.nih.gov/tools/primer-blast/>) by using Xm codes from Vector Base. Serial dilutions of cDNA were used for standard curve production to determine PCR efficiency and specificity of the primer pairs. The qPCR reactions were performed using the Agilent MXPro Real-Time PCR detection system (Agilent Technologies, Stratagene, USA).

A total volume of 20 μl contained 10 μl Brilliant III SYBR Green, 300 nM of primers, 2.5 μl of cDNA (1:10 dilution) and the total volume made up with sterile-distilled water. The cycling conditions for all primer sets consisted of 95°C for 15 minutes followed by 40 cycles of 60°C for 10s, 95°C for 30s, 95°C for 1 minute, 55°C for 30s and 95°C for 30s. Three biological replicates were run for each sample on a plate. Each gene was measured in at least six real-time PCR runs (replicate reactions/biological repeats). The real-time PCR (qPCR) reactions were run along with two normalising 'control' genes, ribosomal S7 and elongation factor EF1. Primer pairs exhibiting a linear relationship between Ct values and template concentration in standard curves, and high PCR amplification efficiency were chosen for further analysis.

Data analysis of gene expression

Outlier data points were identified and excluded from analysis based on obvious deviations in both the normal shape of amplification curves and the Ct values of other repeated observations (biological triplicates). The mean of the threshold cycles (Ct) for each gene were normalised against the average values for ribosomal S7 and elongation factor EF1. To compare gene expression between treatments, we calculated the $\Delta\Delta Ct$ [51]. The

fold-change in gene expression for each target gene, normalized to the ribosomal *S7* and elongation factor *EF1* relative to a susceptible strain of *An. arabiensis* from Dar es Salaam, was calculated according to the $2^{-\Delta\Delta Ct}$ method incorporating PCR efficiency [51]. Basic data analysis (regression and t-tests were performed in Excel. with $p < 0.05$ used to assess significant difference between treatments for the t-tests.

Ethical considerations

The study received ethical clearance from the KCMC Research Ethics Committee.

Results

Microarray experiments involving Moshi samples indicated that 87 probes were significantly up regulated in all three comparisons and 85 down-regulated coming from a total of 133 genes (Figure 2). There was a high correlation among fold changes between the different experimental comparisons, suggesting consistency of results (Figure 3). Surprisingly, the genes that were down regulated include multiple detoxification genes, including several that have come up as top upregulated hits in previous *An. gambiae* s.s. arrays, including P450s repeatedly resistance-associated in previous work on *An. gambiae* s.s. (*CYP6M2* and *CYP6P3*). Among the up-regulated genes were 3 independent ABC transporters, one of which *ABCB4* (represented by 3 alternate transcripts) exhibited almost all significant probes. Also 3 probes for *GSTe7* were up-regulated. Full microarray results from both experiments have been submitted as Additional file 1. However, few genes were selected for qPCR. Of about thirty primers that were designed, only six passed the criteria for qPCR. These primers are shown in Table 1 below including the primers for

CYP4G16, the gene that was documented in other studies reporting on insecticide resistance in *An. arabiensis* and two normalising 'control' genes, ribosomal *S7* and elongation factor *EF1* (Table 1). Only two genes, *ABC 2060* and *CYP4G16* were the most significantly over-transcribed genes in qPCR (Figure 4).

Discussion

Transcriptional analysis of pyrethroid resistance in the four day old Lower Moshi *An. arabiensis* females was investigated using the *An. gambiae* whole genome microarray [25] and quantitative real-time qPCR. Since a fully sequenced genome for *An. arabiensis* was not available; we used heterologous hybridization of the *An. arabiensis* samples to an *An. gambiae* microarray. The microarray analysis of this study clearly indicates that the differences among arrays could not be appropriately determined. In the previous studies, the use of *An. gambiae* 'detox chip' could successfully determine the gene expression of pyrethroid resistant populations of other species including *An. funestus* [53], *An. stephensi* [54] and *An. arabiensis* [28]. Gene expression of pyrethroid resistant *An. arabiensis* populations from Pemba Island was also successfully determined by using *An. gambiae* whole genome microarray [40]. The microarray analysis of this study clearly indicates that there were some very interesting hits among the up regulated genes most notably three independent ABC transporters and *GSTe7*.

Transcription of four genes was evaluated in this study using quantitative real-time (qPCR). It included one ABC transporter gene *ABC 2060* that was up regulated, two P450s (*CYP6M2*, *CYP6P3*) that were down regulated in microarray experiments but previously associated with pyrethroid resistance and one P450 gene *CYP4G16* that was identified through published literature and implicated





Figure 2 Volcano plot of all probes significant in all of three experimental comparisons of *An. arabiensis* permethrin-exposed survivors vs. unexposed samples (average of the three experiments). Red triangles show probes of candidate genes chosen for subsequent qRT-PCR, with probes for the same gene within dashed ovals; the tight cluster of overexpressed gene probes are grouped together.



in permethrin resistance in *An. arabiensis* [40]. Only two genes, CYP4G16 and ABC 2060, showed high expression levels ($p < 0.05$) in the pyrethroid resistant Lower Moshi *An. arabiensis* and it is possible that the two genes may be playing a role in the observed resistance to pyrethroids in the Lower Moshi *An. arabiensis* population. The low levels of expression of the two P450s, CYP6M2 and CYP6P3 in the pyrethroid resistant Lower Moshi *An. arabiensis* implies that the observed pyrethroid resistance results from other mechanisms rather than insecticide metabolism.

CYP4G16 has previously been recorded as an over-transcribed gene in *An. arabiensis* strain from Cameroon [28], Pemba Island Zanzibar [40] and in a laboratory strain of *An. arabiensis* from Sudan that was resistant to DDT, permethrin and deltamethrin [41]. In Pemba Island up-

regulation of CYP4G16 in resistant *An. arabiensis* ranged from 2.0 to 4.5 fold compared to susceptible strains of *An. arabiensis* from Dar es Salaam and Unguja and it has been suggested that it plays a role in cuticular-based resistance. Cytochrome P450s CYP4G17 and CYP4G16 of the *Anopheles gambiae* are closely related to the *Drosophila melanogaster* CYP4G1, the specific insect P450 with decarboxylase function that is involved in hydrocarbon biosynthesis. CYP4G16 and CYP4G17 proteins are found in the mosquito head and the carcass of the abdomen, MozAtlas [55] but not the midgut or malpighian tubules, where P450 detoxification genes have been localized previously.

In this study, metabolic resistance to pyrethroids in Lower Moshi *An. arabiensis* has been shown to be associated with increased transcription of P450 gene, CYP4G16 and ABC transporter. Following qPCR analysis, CYP4G16

Table 1 Primers used in quantitative real-time PCR (qPCR) (F = forward, R = reverse)

Primer	Primer sequence	Fragment length (bp)	Reference
ABC 2060	F: 5'-AATGCACTGCTTTCGGA-3' R: 5'-GACCATCCCACTGTTTCGGA-3'	90	This paper
CYP6P3	F: 5'-GTGATTGACGAAACCCTTCGGAAGT-3' R: 5'-GCACCAGTGTTCGTTTCGGA-3'	N/A	Witzig et al., 2013 [52]
CYP6M2	F: 5'-TACGATGACAACAAGGGCAAG-3' R: 5'-GCGATCGTGAAGTACTGG-3'	N/A	Witzig et al., 2013 [52]
CYP4G16	F: 5'-AGCTGAACGGATACCTGGACCGA-3' R: 5'-AACACGGAGTGCGAACTGCCAAC-3'	81	Jones et al., 2013 [40]
S7	F: 5'-AGAACCAGCAGACCACCATC-3' R: 5'-GCTGCAAACCTTCGGCTATTC-3'	149	Jones et al., 2013 [40]
EF	F: 5'-GGCAAGAGGCATAACGATCAATGCG-3' R: 5'-GTCCATCTGCGACGCTCCGG-3'	N/A	Jones et al., 2013 [40]




Figure 4 Relative expression levels of four candidate genes between sample groups of *An. arabiensis* differing in predicted susceptibility. Data are delta-delta CT values calculated from 5–9 biological repeats. **P < 0.01; *P < 0.05 (t-tests).

was over-expressed in Lower Moshi pyrethroid resistant *An. arabiensis* compared to susceptible strains of *An. arabiensis* from Dar es Salaam. (FC = 3.92, $p = 0.007$). The gene ABC 2060 was over-expressed with the FC of 5.07, ($p = 0.009$). ABC transporters have been linked to insecticide resistance in several species such as *Helicoverpa armigera* [56], *Aedes spp.* [57,58], *Culex pipiens* complex [59] and *Anopheles gambiae* [60]. ABC transporters have also been correlated with insect resistance to *Bacillus thuringiensis* [61]. Members of the ABC-transporter family (ABC-transporters) pump foreign molecules out of insect cells using an ATP-dependent mechanism [62–65]. They are also involved in lipid transport to the cuticle [62]. Interestingly, AGAP002060 is a member of the ABCCH family, which have been implicated in transportation of lipids to the cuticle [62]. The previous study on *An.arabiensis* populations a few kilometers from Mabogini village [45] (the current study site) revealed over-transcription of esterase AGAP006227, the UDPGT AGAP006775 and the cytochrome P450s CYP9J4 and CYP6P1. However, its involvement in insecticide resistance is not clear. In addition, the gene AGAP000987 encoding the cuticle protein CPAP3-A1b was also strongly over-transcribed in *An. arabiensis* in the area. Cuticle proteins play an important role in pyrethroid resistance through altered insecticide penetration and over-transcription of genes encoding them have been reported in pyrethroid resistant species including *Anopheles species* [45,66].

Conclusions

The study compared the gene expression profiles of permethrin-resistant and susceptible populations of *An.*

arabiensis using quantitative real-time PCR (qPCR). *CYP6P3* is one of the leading candidate genes for metabolic resistance in West African *An. gambiae* and its down regulation suggests that quite different mechanisms may operate in the different species.

Based on the quantitative real-time PCR q(PCR), one P450 monooxygenase gene *CYP4G16* and ABC transporter ABC 2060 were over-transcribed in permethrin resistance in Lower Moshi *An. arabiensis*. However, further investigation is needed to clarify its role in observed resistance. There is no real evidence that ABC 2060 transporter gene is linked to observed resistance, although significant lower expression of this gene was observed in a susceptible strain from Dar es Salaam than the Moshi population (i.e. either survivors or Moshi control). However, the two genes identified are most likely not the only genes involved in pyrethroid resistant *An. arabiensis* from Lower Moshi. Further investigation is needed especially in microarray analysis to identify more genes that may be involved in insecticide resistance in the country and its effects on the current malaria control interventions.

Additional file

Additional file 1: Full microarray results from both experiments.

Abbreviations

ITNs: Insecticide-treated nets; LLINs: Long-lasting insecticide treated nets; IRS: Indoor residual spraying; DDT: Dichlorodiphenyltrichloroethane; OPs: Organophosphates; GSTs: Glutathione S-transferases; P450: Cytochrome P450 monooxygenase enzymes; RNA: Ribonucleic acid; qPCR: Quantitative Polymerase Chain Reaction; RT-qPCR: Reverse transcription and quantitative polymerase chain reaction; cDNA: Complementary Deoxyribonucleic acid;

CYP: Genes encoding cytochrome; NCBI: National Center for Biotechnology Information; ANOVA: Analysis of variance; MAANOVA: Multivariate Analysis of Variance; MCDCC: Malaria Capacity Development Consortium; KCMC: Kilimanjaro Christian Medical Centre; WHO: World Health Organization.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

JM conceived, designed, and performed the experiments; participated in data analysis and interpretation; wrote the first draft of the manuscript. CMJ participated in RNA extractions, labelling, hybridizations and scanning of the microarrays; contributed to data analysis and interpretation and reviewed the manuscript. BK contributed to sampling, processing and preservation of samples and review of the manuscript. KS processed mosquitoes and performed reverse transcription and qPCR. FM and MR critically revised the manuscript. HR provided technical support and facilitated the implementation of the microarrays work. DW co-designed the study, provided technical support, contributed to data analysis and interpretation and critically reviewed the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

We are grateful to Hyacinthe Toe, a PhD student at Liverpool School of Tropical Medicine for orienting JM in the laboratory for microarray analysis. Our sincere gratitude to the Vector group of Liverpool School of Tropical Medicine for their support and encouragement during microarray analysis; Professor Martin Donnelly conceived the loop design for microarray analysis. We sincerely express our special thanks to Mr. Augustine Mtui, Rashid Athumani and Issa Masenga for the field larval collections and rearing; Grace Chuwa and Dickson Matowo for mosquito processing and preservation. This work was supported by the Malaria Capacity Development Consortium (MCDCC) which is funded by The Wellcome Trust (Grant number WT084289MA).

Author details

¹Kilimanjaro Christian Medical University College (KCMUCo), Moshi, Tanzania. ²Liverpool School of Tropical Medicine (LSTM), Liverpool, UK. ³London School of Hygiene and Tropical Medicine (LSHTM), London, UK. ⁴Pan-African Malaria Vector Control Consortium (PAMVERC), Moshi, Tanzania. ⁵National Institute for Medical Research (NIMR), Tukuyu, Tanzania.

Received: 23 April 2014 Accepted: 15 June 2014

Published: 19 June 2014

References

- Zaim M, Jambulingam P: *Global insecticide use for vector-borne disease control 3rd edition*. Geneva: World Health Organization; 2007.
- WHO: *World malaria report 2013*. Geneva: World Health Organization; 2013.
- Abuelmaali SA, Elaagip AH, Basheer MA, Frah EA, Ahmed FTA, Elhaj HFA, Seidahmed OME, Hamid MMA WD: **Impacts of agricultural practices on insecticide resistance in the malaria vector *Anopheles arabiensis* in Khartoum state, Sudan.** *PLoS One* 2013, **8**(12). doi:10.1371/annotation/d40e811c-993d-40ec-8294-420402282448.
- Czeher C, Labbo R, Arzika I, Duchemin JB: **Evidence of increasing Leu-Phe knock down resistance- mutation in *Anopheles gambiae* from Niger following a nationwide long-lasting insecticide treated nets implementation.** *Malar J* 2008, **7**:189.
- Protopopoff N, Verhaegen K, Van Bortel W, Roelants P, Marcotty T, Baza D, D'Alessandro U, Coosemans M: **A significant increase in *kdr* in *Anopheles gambiae* associated with an intensive vector control intervention in Burundi highlands.** *Trop Med Int Health* 2008, **13**:1479-1487.
- Yadoulton A, Martin T, Padonou G, Chandre F, Asidi A, Djogbenou L, Dabiré R, Aikpon R, Boko M, Gliitho I, Akogbeto M: **Cotton pest management practices and the selection of pyrethroid resistance in *Anopheles gambiae* population in Northern Benin.** *Parasit Vectors* 2011, **4**:60.
- Diabate A, Baldet T, Chandre F, Akogbeto M, Guiguemde RT, Darriet F, Brengues C, Guillet P, Hemingway J, Graham JS, Hougard JM: **The role of agricultural use of insecticides in resistance to pyrethroids in *Anopheles gambiae* s.l in Burkina Faso.** *Am J Trop Med Hyg* 2002, **67**:617-622.
- Ranson H, N'Guessan R, Lines J, Moiroux N, Nkuni Z, Corbel V: **Pyrethroid resistance in African anopheline mosquitoes: what are the implications for malaria control?** *Trends Parasitol* 2011, **27**(2):91-98.
- Zaim M, Aitio A, Nakashima N: **Safety of pyrethroid-treated mosquito nets.** *Med Vet Entomol* 2000, **14**:1-5.
- Global Malaria Programme WHO: *Global plan on insecticide resistance management in malaria vectors (GPIRM)*. Geneva: World Health Organization; 2012.
- Coleman M, Hemingway J: **Insecticide resistance monitoring and evaluation in disease transmitting mosquitoes.** *J Pest Sci* 2007, **32**(2):69-76.
- Hemingway J, Hawkes NJ, McCarroll L, Ranson H: **The molecular basis of insecticide resistance in mosquitoes.** *Insect Biochem Mol Biol* 2004, **34**(7):653-665.
- Martinez Torres D, Chandre F, Williamson MS, Darriet F, Berge JB, Devonshire AL, Guillet P, Pasteur N, Pauron D: **Molecular characterization of pyrethroid knockdown resistance (*kdr*) in the major malaria vector *Anopheles gambiae* s.s.** *Insect Mol Biol* 1998, **7**(2):179-184.
- Ranson H, Jensen B, Vulule JM, Wang X, Hemingway J, Collins FH: **Identification of a point mutation in the voltage-gated sodium channel gene of Kenyan *Anopheles gambiae* associated with resistance to DDT and pyrethroids.** *Insect Mol Biol* 2000, **9**(5):491-497.
- Ranson H, Rossiter L, Orтели F, Jensen B, Wang XL, Roth CW, Collins FH, Hemingway J: **Identification of a novel class of insect glutathione S-transferases involved in resistance to DDT in the malaria vector *Anopheles gambiae*.** *Biochem J* 2001, **359**:295-304.
- Gunasekaran K, Muthukumaravel S, Sahu SS, Vijayakumar T, Jambulingam P: **Glutathione S-transferase activity in Indian vectors of malaria: A defense mechanism against DDT.** *J Med Entomol* 2011, **48**(3):561-569.
- Prapanthadara LA, Ketterman AJ: **Qualitative and quantitative changes in glutathione S-transferases in the mosquito *Anopheles gambiae* confer DDT-resistance.** *Biochem Soc Trans* 1993, **21**(3):304S.
- Vontas JG, Small GJ, Hemingway J: **Glutathione S-transferases as antioxidant defence agents confer pyrethroid resistance in *Nilaparvata lugens*.** *Biochem J* 2001, **357**:65-72.
- Kostaropoulos I, Papadopoulos AI, Metaxakis A, Boukouvala E, Papadopoulou-Mourkidou E: **Glutathione S-transferase in the defence against pyrethroids in insects.** *Insect Biochem Mol Biol* 2001, **31**(4-5):313-319.
- Riveron JM, Yunta C, Ibrahim SS, Djouaka R, Irving H, Menze BD, Ismail HM, Hemingway J, Ranson H, Albert A, Wondji CS: **A single mutation in the *GSTe2* gene allows tracking of metabolically based insecticide resistance in a major malaria vector.** *Gen Biol* 2014, **15**:R27.
- Herath PR, Miles SJ, Davidson G: **Fenitrothion (OMS 43) resistance in the taxon *Anopheles culicifacies* Giles.** *J Trop Med Hyg* 1981, **84**(2):87-88.
- Hemingway J: **The genetics of malathion resistance in *Anopheles stephensi* from Pakistan.** *Trans R Soc Trop Med Hyg* 1983, **77**(1):106-108.
- Feyerisen R, Lawrence IG, Kostas I, Sarjeet SG: **Insect Cytochrome P450.** In *Comprehensive Molecular Insect Science*. Amsterdam: Elsevier; 2005:1-77.
- Muller P, Warr E, Stevenson BJ, Pignatelli PM, Morgan JC, Stephen A, Yawson E, Mitchell SN, Ranson H, Hemingway J, Paine MJ, Donnelly MJ: **Field-caught permethrin-resistant *Anopheles gambiae* overexpress CYP6P3, a P450 that metabolises pyrethroids.** *PLoS Genet* 2008, **4**:e000286.
- Djouaka RF, Bakare AA, Coulibaly ON, Akogbeto MC, Ranson H, Hemingway J, Strode C: **Expression of the cytochrome P450s, CYP6P3 and CYP6M2 are significantly elevated in multiple pyrethroid resistant populations of *Anopheles gambiae*s. from Southern Benin and Nigeria.** *BMC Genomics* 2008, **9**:538.
- Muller P, Donnelly MJ, Ranson H: **Transcription profiling of a recently colonised pyrethroid resistant *Anopheles gambiae* strain from Ghana.** *BMC Genomics* 2007, **8**:3626.
- Stevenson BJ, Bibby J, Pignatelli P, Muangnoicharoen S, O'Neill PM, Lian LY, Muller P, Nikou D, Steven A, Hemingway J, Sutcliffe MJ, Paine MJ: **Cytochrome P450 6 M2 from the malaria vector *Anopheles gambiae* metabolizes pyrethroids: Sequential metabolism of deltamethrin revealed.** *Insect Biochem Mol Biol* 2011, **41**(7):492-502.
- Muller P, Chouaibou M, Pignatelli P, Etang J, Walker ED, Donnelly MJ, Simard F, Ranson H: **Pyrethroid tolerance is associated with elevated expression of antioxidants and agricultural practice in *Anopheles arabiensis* sampled from an area of cotton fields in Northern Cameroon.** *Mol Ecol* 2008, **17**(4):1145-1155.
- Mitchell SN, Stevenson BJ, Muller P, Wilding CS, Egyir-Yawson A, Field SG, Hemingway J, Paine MJ, Ranson H, Donnelly MJ: **Identification and validation of a gene causing cross-resistance between insecticide classes in *Anopheles gambiae* from Ghana.** *Proc Natl Acad Sci U S A* 2012, **109**(16):6147-6152.

30. Gillies MT, Coetzee M: **A supplement to the Anophelinae of Africa south of the Sahara (Afrotropical region).** *Johannesburg S Afr Inst Med Res* 1987, **55**:1–143
31. Coetzee M, Crag M, Le Sueur D: **Distribution of African malaria mosquitoes belonging to the *Anopheles gambiae* complex.** *Parasitol Today* 2000, **16**:74–77.
32. Minakawa N, Sonye G, Mogi M, Githeko A, Yan G: **The effects of climatic factors on the distribution and abundance of malaria vectors in Kenya.** *J Med Entomol* 2002, **39**:833–841.
33. Verhaeghen K, Bortel WW, Roelants P, Backeljau T, Coosemans M: **Detection of the East and West African *kdr* mutation in *Anopheles gambiae* and *Anopheles arabiensis* from Uganda using a new assay based on FRET/Melt Curve analysis.** *Malar J* 2006, **5**:16.
34. Himeidan YE, Chen H, Chandre F, Donnelly MJ: **Short report: permethrin and DDT resistance in the malaria vector *Anopheles arabiensis* from eastern Sudan.** *Am J Trop Med Hyg* 2007, **77**:1066–1068.
35. Abdalla H, Matambo TS, Koekemoer LL, Mnzava AP, Hunt RH, Coetzee M: **Insecticide susceptibility and vector status of natural populations of *Anopheles arabiensis* from Sudan.** *Trans R Soc Trop Med Hyg* 2008, **102**:263–271.
36. Ndjemai HNM, Patchoké S, Atangana J, Etang J, Simard F, Bilong Bilong CF, Reimer L, Cornel A, Lanzaro CG, Fondjo E: **The distribution of insecticide resistance in *Anopheles gambiae* s.l. populations from Cameroon: an update.** *Trans R Soc Trop Med Hyg* 2008, **103**(11):1127–1138.
37. Kulkarni MA, Rowland M, Alifrangis M, Moshia FW, Matowo J, Malima R, Justin P, Kweka E, Lyimo I, Magesa S, Salanti A, Rau ME, Drakeley C: **Occurrence of leucine to phenylalanine knockdown resistance (*kdr*) mutation in *Anopheles arabiensis* populations Tanzania detected by a high throughput SSOP-ELISA method.** *Malar J* 2006, **5**:56.
38. Protopopoff N, Matowo J, Malima R, Kavishe R, Kaaya R, Wright A, West P, Kisinza W, Moshia FW, Kleinschmidt I, Rowland M: **High level of resistance to pyrethroids and DDT and reduced susceptibility to bendiocarb in malaria vector *Anopheles gambiae* in North Western Tanzania.** *Malar J* 2013, **12**:149.
39. Matowo J, Kitau J, Kabula B, Oxborough RM, Moshia FW, Rowland M: **Dynamics of pyrethroid resistance and the frequency of *kdr* mutation in *Anopheles arabiensis* in rural villages of Lower Moshi, North-Eastern Tanzania.** *J Parasitol Vector Biol* 2014, **6**(3):31–41.
40. Jones CM, Haji KA, Khatib BO, Bagi J, Mcha J, Devine GJ, Daley M, Kabula B, Ali AS, Majambere S, Ranson H: **The dynamics of pyrethroid resistance in *Anopheles arabiensis* from Zanzibar and an assessment of the underlying genetic basis.** *Parasite Vectors* 2013, **6**:343.
41. Nardini L, Christian RN, Coetzer N, Ranson H, Coetzee M, Koekemoer LL: **Detoxification enzymes associated with insecticide resistance in laboratory strains of *Anopheles arabiensis* of different geographic origin.** *Parasite Vectors* 2012, **5**:113.
42. Mouatcho JC, Munhenga G, Hargreaves K, Brooke BD, Coetzee M, Koekemoer LL: **Pyrethroid resistance in a major African malaria vector *Anopheles arabiensis* from Mamfene, northern KwaZulu-Natal, South Africa.** *S Afr J Sci* 2009, **105**:127–131.
43. **Universal Coverage Campaign - Final Impact.** 2011. http://medatanzania.org/?p=ucc_final.
44. Matowo J, Kulkarni MA, Moshia FW, Oxborough R, Kitau J, Tenu F, Rowland M: **Biochemical basis of permethrin resistance in *Anopheles arabiensis* from Lower Moshi, north-eastern Tanzania.** *Malar J* 2010, **9**:193.
45. Nkya TE, Akhouayri I, Kisinza W, David JP: **Impact of environment on mosquito response to pyrethroid insecticides: facts, evidences and prospects.** *Insect Biochem Mol Biol* 2013, **43**:407–416.
46. Rogers DW, Whitten MMA, Thailayil J, Soichot J, Levashina EA, Catteruccia F: **Molecular and cellular components of the mating machinery in *Anopheles gambiae* females.** *Proc Natl Acad Sci U S A* 2008, **105**(49):19390–19395.
47. WHO: **Test procedures for insecticide resistance monitoring in malaria vector mosquitoes.** Geneva: World Health Organization; 2013.
48. Scott JA, Brogdon WG, Collins FH: **Identification of single specimens of the *Anopheles gambiae* complex by the polymerase chain reaction.** *Am J Trop Med Hyg* 1993, **49**:520–529.
49. Smyth GK L: **Linear models for microarray data.** In *Bioinformatics and Computational Biology Solutions using R and BIOCONDUCTOR*. Edited by Gentleman R, Carey Y, Dudoit S, Irizarry R, Huber W. New York: Springer; 2005:397–420.
50. Wu H, Kerr M, Cui X, Churchill G: **MAANOVA: A Software Package for the Analysis of Spotted cDNA Microarray Experiments.** In *The Analysis of Gene Expression Data*. Edited by Parmigiani G, Garrett E, Irizarry R, Zeger S. London: Springer; 2003:313–341.
51. Schmittgen TD, Livak KJ: **Analyzing real-time PCR data by the comparative C-T method.** *Nat Protoc* 2008, **3**:1101–1108.
52. Witzig C, Parry M, Morgan JC, Irving H, Steven A, Cuamba N, Kerah-Hinzoumbé C, Ranson H, Wondji CS: **Genetic mapping identifies a major locus spanning P450 clusters associated with pyrethroid resistance in *kdr*-free *Anopheles arabiensis* from Chad.** *Heredity* 2013, **110**:389–397.
53. Christian RN, Strode C, Ranson H, Coetzer N, Coetzee M, Koekemoer L: **Microarray analysis of a pyrethroid resistant African malaria vector, *Anopheles funestus*, from Southern Africa.** *Pestic Biochem Physiol* 2011, **99**:140–147.
54. Vontas J, David J-P, Nikou D, Hemingway J, Christophides GK, Louis C, Ranson H: **Transcriptional analysis of insecticide resistance in *Anopheles stephensi* using cross-species microarray hybridization.** *Insect Mol Biol* 2007, **16**:315–324.
55. Baker D, Nolan T, Fischer B, Pinder A, Crisanti A, Russell S: **A comprehensive gene expression atlas of sex- and tissue-specificity in the malaria vector, *Anopheles gambiae*.** *BMC Genomics* 2011, **12**(1):296.
56. Aurade RM, Jayalakshmi SK, Sreeramulu K: **P-glycoprotein ATPase from the resistant pest, *Helicoverpa armigera*: purification, characterization and effect of various insecticides on its transport function.** *Biochim Biophys Acta* 2010, **1798**:1135–1143.
57. Bariami V, Jones CM, Poupardin R, Vontas J, Ranson H: **Gene Amplification, ABC Transporters and Cytochrome P450s: Unraveling the Molecular Basis of Pyrethroid Resistance in the Dengue Vector, *Aedes aegypti* PLoS Negl Trop Dis** 2012, **6**(6):e1692. doi:10.1371/journal.pntd.0001692.
58. Porretta D, Gargani M, Bellini R, Medici A, Punelli F, Urbanelli S: **Defence mechanisms against insecticides temephos and diflubenzuron in the mosquito *Aedes caspius*: the P-glycoprotein efflux pumps.** *Med Vet Entomol* 2008, **22**:48–54.
59. Buss DS, McCaffery AR, Callaghan A: **Evidence for p-glycoprotein modification of insecticide toxicity in mosquitoes of the *Culex pipiens* complex.** *Med Vet Entomol* 2002, **16**:218–222.
60. Roth CW, Holm I, Graille M, Dehoux P, Rzhetsky A, Wincker P, Weissenbach J, Brey PT: **Identification of the *Anopheles gambiae* ATP-binding cassette transporter superfamily genes.** *Mol Cells* 2003, **15**:150–158.
61. Gahan LJ, Pauchet Y, Vogel H, Heckel DG: **An ABC transporter mutation is correlated with insect resistance to *Bacillus thuringiensis* Cry1Ac toxin.** *PLoS Genet* 2010, **6**:e1001248.
62. Broehan G, Kroeger T, Lorenzen M, Merzendorfer H: **Functional analysis of the ATP-binding cassette (ABC) transporter gene family of *Tribolium castaneum*.** *BMC Gen* 2013, **14**:6.
63. Higgins CF, Linton KJ: **The ATP switch model for ABC transporters.** *Nat Struct Mol Biol* 2004, **11**:918–926.
64. Linton KJ, Higgins CF: **Structure and function of ABC transporters: the ATP switch provides flexible control.** *PLoS Arch* 2007, **453**:555–567.
65. George AM, Jones PM: **Perspectives on the structure-function of ABC transporters: the Switch and Constant Contact Models.** *Prog Biophys Mol Biol* 2012, **109**:95–107.
66. Wood OR, Hanrahan S, Coetzee M, Koekemoer LL, Brooke DB: **Cuticle thickening associated with pyrethroid resistance in the major malaria vector *Anopheles funestus*.** *Parasite Vectors* 2010, **3**:67.

doi:10.1186/1756-3305-7-274

Cite this article as: Matowo et al.: Genetic basis of pyrethroid resistance in a population of *Anopheles arabiensis*, the primary malaria vector in Lower Moshi, north-eastern Tanzania. *Parasites & Vectors* 2014 **7**:274.