

POSTER PRESENTATION

Open Access

Anopheles maculipennis complex-responsible for the re-emergence of malaria in Romania?

ML Ivănescu*, D Acatrinei, L Miron

From The 1st Conference on Neglected Vectors and Vector-Borne Diseases (EurNegVec): with Management Committee and Working Group Meetings of the COST Action TD1303 Cluj-Napoca, Romania. 8-11 April 2014

Between 2010 and 2012, in Iași City of Romania 5 species belonging to the complex *Anopheles maculipennis* were identified by PCR: *A. melanoon*, *A. labranchiae*, *A. atroparvus*, *A. messeae* and *A. maculipennis* sp. The last three species were involved in the transmission of malaria in Romania between 1892 and 1961. The species *A. labranchiae* was signalled for the first time in Romania, being considered as the main vector of malaria in Europe. The identified samples of *Anopheles labranchiae* were two stage IV larvae, concluding with the adaptation of the species to the climate of Iași City. Using a new mathematical model realized and implemented by ourselves, based on the construction of a function of interpolation of Lagrange polynomial type, we realized an extrapolation of the evolution of temperatures for the year 2030, demonstrating the existence of favourable conditions of development of both the vector and the malaria parasite. Therefore, it was a temperature raise by 0.72°C in the whole country, as compared to the period of malaria eradication in Romania, and the extrapolation of the evolution of temperatures in the year 2030 showed a raise by 0.8°C, result which coincides with the prognosis made by prestigious research institutes: NIES, CSIRO, HCCPR, MPIM and NCAR, which show a raise by 0.8-1.7° in 2030. In Iași City, the prognosis made for 2030 shows an average of 24°C for this year in the spring-summer months, temperature which is ideal for the development of the life cycle of mosquitoes, but also for the development of the parasite inside the vector.

Taking into account the fact that Iasi, as a cultural city, hosts many foreign students who are carriers of *Plasmodium*, coming from malaria endemic areas, and the immigration of the population for work in countries

of the African continent, the human reservoir of *Plasmodium* will be permanently maintained. Correlating the three factors implicated in the transmission of malaria: favourable environmental conditions, presence of the vector and of the human reservoir, we could highlight the growing risk of the re-emergence of malaria in Romania.

Published: 1 April 2014

doi:10.1186/1756-3305-7-S1-P15

Cite this article as: Ivănescu et al.: *Anopheles maculipennis* complex-responsible for the re-emergence of malaria in Romania? *Parasites & Vectors* 2014 **7**(Suppl 1):P15.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: larissa81@yahoo.com

Department of Clinics, Faculty of Veterinary Medicine, "Ion Ionescu de la Brad" University of Agricultural Sciences and Veterinary Medicine, Iasi, Romania