

RESEARCH

Open Access



# Analyses of the expression, immunohistochemical properties and serodiagnostic potential of *Schistosoma japonicum* peroxiredoxin-4

Minh-Anh Dang-Trinh<sup>1,2,3</sup>, Jose Ma. M. Angeles<sup>4</sup>, Kharleezelle J. Moendeg<sup>5</sup>, Adrian Miki C. Macalanda<sup>6</sup>, Thu-Thuy Nguyen<sup>1</sup>, Luna Higuchi<sup>1</sup>, Shotaro Nakagun<sup>7</sup>, Masashi Kirinoki<sup>8</sup>, Yuichi Chigusa<sup>8</sup>, Yasuyuki Goto<sup>9</sup> and Shin-ichiro Kawazu<sup>1,2\*</sup>

## Abstract

**Background:** *Schistosoma japonicum*, which inhabits the mesenteric vein of the mammalian hosts for about 20 to 30 years, is subjected to the oxidative stresses from the host defense mechanism during their intra-mammalian stages. To counteract this host immune attack, the parasite utilizes their antioxidant system for survival inside the host. Peroxiredoxins (Prxs), thiol-specific antioxidant proteins, play an essential role for protecting the parasite against oxidative stress by reducing hydrogen peroxide to water. Only three types of 2-Cys Prxs have been previously characterized in *S. japonicum* whereas a fourth Prx has been identified for *Schistosoma mansoni* as Prx-4. A sequence coding homologous to this gene in the *S. japonicum* database was identified, characterized and expressed as recombinant SjPrx-4 protein (rSjPrx-4). Furthermore, rSjPrx-4 was evaluated in this study for its diagnostic potentials in detecting *S. japonicum* infection in humans.

**Results:** The gene found in the parasite genome contained 2 active-site cysteines with conserved sequences in the predicted amino acid (AA) sequence and showed 75% identity with that of the previously characterized Prx (TPx-1) of *S. japonicum*. The gene was expressed in different stages of schistosome life-cycle with highest transcription level in the adult male. The gene was cloned into a plasmid vector and then transfected into *Escherichia coli* for expression of rSjPrx-4. Anti-rSjPrx-4 mouse sera recognized native SjPrx-4 in egg and adult worm lysate by western blotting. The result of a mixed function oxidation assay in which rSjPrx-4 prevented the nicking of DNA from hydroxyl radicals confirmed its antioxidant activity. Subsequently, immunolocalization analysis showed the localization of SjPrx-4 inside the egg, on the tegument and in the parenchyma of the adult worm. Enzyme-linked immunosorbent assay results showed that rSjPrx-4 has 83.3% sensitivity and 87.8% specificity. Its diagnostic potential was further evaluated in combination with recombinant SjTPx-1 protein, yielding an improved sensitivity and specificity of 90% and 92.7%, respectively.

**Conclusions:** These results suggest that SjPrx-4 plays a role as an antioxidant dealing with oxidative stresses of *S. japonicum*, and its diagnostic potential improved by coupling it with SjTPx-1 is a proof for developing a serological test with better diagnostic performance for human schistosomiasis.

\*Correspondence: skawazu@obihiro.ac.jp

<sup>1</sup> National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Hokkaido, Japan  
Full list of author information is available at the end of the article



© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

**Keywords:** *Schistosoma japonicum*, Biomarker, Peroxiredoxin-4, ELISA, Diagnosis

## Background

Schistosomiasis japonica, a zoonotic disease caused by *Schistosoma japonicum*, still remains as an economic and public health concern in China, the Philippines and some parts of Indonesia. A major consideration making the disease prevalence less controllable is the use of diagnostic methods with low efficiency in detecting infection and assessing the success of control measures [1]. The standard test for the diagnosis of intestinal schistosomiasis in endemic areas is still parasitological detection of eggs in stool samples such as Kato-Katz [2]. Although the Kato-Katz is simple and has a specificity of 100%, it is time-consuming, laborious, and poses unsatisfactory sensitivity mostly in endemic areas with low prevalence of schistosomiasis [3]. Therefore, many individuals with light to moderate intensity of infection were underdiagnosed using Kato-Katz method [4, 5].

Immunodiagnostic tools based on schistosome antigens have a high sensitivity and are less time consuming as compared to coprological diagnostic tools. Thus, these assays have become an attractive alternate for schistosomiasis surveillance especially in the locations where the disease has been nearly eliminated. Many antigens with high sensitivity and specificity have been reported as promising markers for diagnosing *S. japonicum* infection [6, 7]. However, single recombinant antigens used for the serodiagnosis of human schistosomiasis were found to have less sensitivity with lower *S. japonicum* infection intensity level in different human cohorts [8]. Employing multivalent cocktail antigens could possibly improve the low sensitivity of single recombinant antigens in leishmaniasis [9], in clonorchiasis [10] and in malaria [11] as well as schistosomiasis [8]. Moreover, a cocktail-ELISA showed the improvement of specificities in humans, water buffaloes, and dogs for schistosomiasis japonica [12]. Therefore, identification of the synergistic diagnostic potentials of different combinations of specific antigens recognized by the host immune system is essential and will help improve the test on detecting the schistosome infection.

The schistosome adult worms can live in the host hepatic mesenteries for about 20–30 years without any treatment being administered to the host [13, 14]. In this environment, the parasite must deal with reactive oxygen species (ROS) mainly produced by the host immune system as well as those produced by the parasite itself during erythrocyte ingestion. To protect themselves, one of the defense mechanisms the parasite uses is through the production of an antioxidant firewall [15].

Peroxiredoxins (Prxs) are members of a family of antioxidant enzymes that detoxify hydrogen peroxide to water [16]. Prxs have been classified into 2-Cys Prx and 1-Cys Prx based on the number of cysteine residues involved in the peroxidase activity [17]. The 2-Cys Prxs are also known as thioredoxin peroxidases (TPx) because they use thiol from the thioredoxin system to carry out peroxidase function [18]. Three types of *S. japonicum* peroxiredoxin (Prx), also called thioredoxin peroxidase (TPx), have been cloned and characterized, namely TPx-1, TPx-2 and TPx-3 [19]. SjTPx-1 was found in the tegumental tissues of the adult worm and in excretory/secretory products [19, 20]. SjTPx-2 was mainly expressed in gut epithelium, vitelline gland, parenchyma and in the sub-tegument of adult worm [19]. SjTPx-3 has mitochondrial signal and was considered to play a role in mitochondria [19, 21]. However, a fourth Prx, reported in *S. mansoni* as SmPrx-4 [22], has not yet been identified in *S. japonicum*.

Moreover, there have been many studies identifying thioredoxin peroxidases as potential antigens for diagnosing various parasitic diseases such as *Fasciola gigantica* infection in water buffaloes [23] and *Plasmodium falciparum* malaria in humans [24]. Among three types of *S. japonicum* thioredoxin peroxidase, SjTPx-2 and SjTPx-3 were evaluated as vaccine candidates against schistosome infection [21, 25] and only SjTPx-1 was identified as a promising candidate antigen for serodiagnosis of *S. japonicum* infection in humans [26], water buffaloes [27] and dogs [28].

In this study, the *S. japonicum* Prx-4 (SjPrx-4) gene was cloned and expressed. Then, its expression levels among different developmental stages and the antioxidant activity of the recombinant SjPrx-4 (rSjPrx-4) were further analyzed. Localization of the protein was also examined and its diagnostic potential for human schistosomiasis was evaluated as a single antigen and cocktail-antigen with SjTPx-1 using enzyme-linked immunosorbent assay (ELISA).

## Methods

### Parasites and animal infection

The *S. japonicum* Yamanashi strain was maintained using the snail intermediate host *Oncomelania hupensis nosophora*. Miracidia were collected after the eggs were hatched in fresh water. Then, *O. h. nosophora* snails were infected with 5 miracidia and were crushed 6 months later to collect the cercariae required in this study. Five female ICR mice aged 5 weeks-old (Clea Inc, Tokyo,

Japan) were used for *S. japonicum* infection. The mice were percutaneously infected with 40 *S. japonicum* cercariae each and sacrificed at 8 weeks post-infection.

#### Human sera

Archived serum samples collected from Filipinos living in Irosin, Sorsogon, Philippines (schistosomiasis endemic areas) in 2015 were used in this study in accordance with the ethical clearance approved by the University of the Philippines, Manila. Thirty schistosomiasis-positive and 41 schistosomiasis-negative serum samples were confirmed by Kato-Katz in triplicate slide examinations of approximately 41.7 mg fecal sample each and stool PCR for *S. japonicum* [26]. Moreover, 45 healthy American sera were used as negative controls (Bioreclamation-IVT, Baltimore, MD, USA). Archived sera from patients infected with *Paragonimus westermani* ( $n = 10$ ), *Entamoeba histolytica* ( $n = 9$ ), *Opisthorchis viverrini* ( $n = 6$ ) and *Trichuris trichiura* ( $n = 1$ ) were also used to test the cross-reactivity. *Entamoeba histolytica*-positive and *T. trichiura*-positive sera were collected from a schistosomiasis-free area in the Philippines and confirmed through microscopic examination. *Opisthorchis viverrini*-positive samples were taken from Thailand and *P. westermani*-positive sera were collected from Japanese patients diagnosed through either clinical manifestations or antibody detection [26].

#### Sequence analysis

The coding sequence of SjPrx-4 (accession no. Sjp\_0053380) was retrieved from GeneDB with the *S. mansoni* Prx-4 sequence as reference using BLASTX in NCBI (<http://www.ncbi.nlm.nih.gov/>). The Compute pI/Mw tool ([http://www.ExPASy.ch/tools/pi\\_tool.html](http://www.ExPASy.ch/tools/pi_tool.html)) was used to calculate the molecular weight and isoelectric point of SjPrx-4. The SignalP 4.0 was used to predict the signal peptide (<http://www.cbs.dtu.dk/services/SignalP/>).

#### Preparation of rSjPrx-4

Primers of SjPrx-4 were designed according to the sequence obtained from GeneDB (accession no. Sjp\_0053380) with restriction enzyme sites of *Bam*HI and *Xho*I (underlined) at the forward (5'-GCCG GAT CCA TGC TGT TAC CAA CG-3') and reverse oligonucleotides (5'-GCCC TCG AGT CAT GTA GAT GAA GAG-3'), respectively. The PCR product was directly cloned into pCR 2.1-TOPO vector (Invitrogen, CA, USA) according to the manufacturer's protocol. And the plasmids were transformed into *Escherichia coli* Mach I (Invitrogen). Selected clones were confirmed by sequencing for the sequence identity. Then the SjPrx-4 gene was digested with *Bam*HI and *Xho*I to be subcloned from pCR 2.1-TOPO vector into pGEX-6P-1 vectors (Novagen,

Madison, USA). Subsequently, pGEX-6P-1 plasmids containing the SjPrx-4 gene were transfected into *E. coli* strain Rosetta (DE3) (Novagen, MA, USA). The expression of rSjPrx-4 was induced by 1 mM isopropyl- $\beta$ -D-thiogalactoside (IPTG) at 25 °C overnight. The bacteria pellets were collected by centrifuging at 10,000 $\times$ g, 4 °C for 20 min and were sonicated in lysis buffer. Cellular debris was pelleted at 8000 $\times$ g, 4 °C for 20 min. The supernatant was collected and purified using glutathione Sepharose 4B resin and cleaved by PreScission Protease to remove glutathione S-transferase (GST)-tag (GE Healthcare BioSciences AB, Uppsala, Sweden) following the manufacturer's instructions. Purified rSjPrx-4 protein was then dialyzed 3 times with phosphate-buffered saline (PBS), pH 7.2. The concentration of the rSjPrx-4 was determined by bicinchoninic acid (BCA) assay (Thermo Fisher Scientific Inc., Rockford, IL, USA), and the protein was stored at -80 °C until use.

#### Anti-rSjPrx-4 serum development

To obtain anti-rSjPrx-4 antibodies, rSjPrx-4 emulsified with the TiterMax Gold<sup>®</sup> adjuvant (Sigma-Aldrich, St. Louis, MO, USA) was inoculated subcutaneously into 8 week-old female ICR mice (100  $\mu$ g rSjPrx-4/200  $\mu$ l inoculum/mouse). Five mice were immunized thrice with a 2-week interval by subcutaneous injection. Then mouse sera were collected, pooled and labeled as anti-rSjPrx-4 mouse serum after the 3rd immunization. The anti-rSjPrx-4 mouse serum was used for immunohistochemistry analysis as primary antibody.

#### Quantitative real time reverse transcription-PCR (qRT-PCR) analysis

The transcription level of the SjPrx-4 gene in each developmental stage was examined by qRT-PCR in triplicates. The intramolluscan stages of *S. japonicum* were obtained from schistosome-infected *O. h. nosophora* whereas the intramammalian stages were from infected mice. The total RNA samples were prepared from adult worm, egg and sporocyst with TRIZOL reagent (Invitrogen, CA, USA) according to the manufacturer's instruction. The PrimeScript RT reagent kit (Takara, Shiga, Japan) was used to synthesize the cDNA library. The forward (5'-GTT ACC AAC GAA ACC GGC G-3') and reverse (5'-CCC AAT CCT CCT GCT TTA CGA-3') primers were used to amplify the SjPrx-4 gene with a size of 273 bp. Triose-phosphate isomerase (TPI) gene (forward primer: 5'-ATG GCA GTA GAG CCG ACA AC-3'; reverse primer: 5'-AAC GCT TAG ACC TTC TGC AA-3') was used as an internal standard gene [19]. Twenty microliters of PCR reaction mixture was prepared containing 0.5  $\mu$ l of template cDNA, 10  $\mu$ l of PowerUp SYBR Green

Master Mix (Applied Biosystems, UAB, Lithuania), 200 nM of each primer, and distilled water up to 20  $\mu$ l. The reactions were run on an Applied Biosystems 7900HT instrument (Applied Biosystems, Foster City, CA, USA) under the following conditions: 95 °C for 3 min, 40 cycles at 95 °C for 30 s and at 60 °C for 1 min, and a dissociation stage as per the instrument guidelines. The  $2^{-\Delta\Delta C_q}$  method [29, 30] was performed to analyze data with TPI gene employed as the internal control. The transcriptional changes of SjPrx-4 gene in the eggs, male and female adult worms were calculated relative to that of the sporocyst. Negative controls not containing the parasite DNA were included in each PCR run.

#### Antioxidant activity of rSjPrx-4

A mixed-function oxidation (MFO) assay was performed to evaluate the antioxidant activity of rSjPrx-4 [31]. The mixture of 50  $\mu$ l containing 25 mM HEPES (pH 7.0), 10 mM dithiothreitol (DTT), 20 mM EDTA and 40  $\mu$ M FeCl<sub>3</sub>, was pre-incubated without or with rSjPrx-4 (10 ng, 100 ng, 1  $\mu$ g and 5  $\mu$ g/ml) for 1 h at 37 °C. After pre-incubation 500 ng of supercoiled pBluescript plasmid DNA (Stratagene, La Jolla, CA, USA) was added into the reaction mixture and incubated for another 3 h at 37 °C. Finally, 0.8% agarose gel electrophoresis was used to assess the nicking of the supercoiled plasmid by MFO. The heat-denatured rSjPrx-4 protein was included as negative control.

#### Western blotting

The specific binding of anti-rSjPrx-4 with SjPrx-4 and the expression of SjPrx-4 in *S. japonicum* eggs and adult worms were determined by western blotting. The egg and adult worm lysate were analyzed on 12% SDS-PAGE and transferred onto a polyvinylidene difluoride (PVDF) membrane. The membrane was blocked with 3% skimmed milk in Tris-buffered saline with 0.05% of Tween 20 (TBS-T) overnight at 4 °C. Then, the membrane was incubated with anti-rSjPrx-4 at 1:100 dilutions in blocking buffer for 1 h at room temperature (RT). After washing 3 times with TBS-T, the membrane was further incubated with an anti-mouse IgG conjugated with horseradish peroxidase (HRP) (GE Healthcare) at dilution of 1:5000 for 1 h at RT. After washing, specific binding of anti-rSjPrx-4 was visualized using SuperSignal HRP chemiluminescent substrates (ThermoFisher Scientific) and an ImageQuant LAS 500 chemiluminescence detection machine (GE Electric, Tokyo, Japan). Membranes containing egg and adult lysate treated with normal mouse serum were used as negative controls, separately.

#### Immunolocalization of SjPrx-4 in schistosomes

Immunohistochemistry was performed on 10% formalin-fixed paraffin embedded sections of adult worms and visceral organs obtained from the infected mice according to standard techniques [32], with some modifications. The heat-induced antigen retrieval method was done with microwave at 97 °C in citrate buffer (pH 6.0) for 15 min. Inhibition of endogenous peroxidase activity was performed by incubating the slides with 0.3% hydrogen peroxide at RT for 10 min. The sections on the slide glasses were incubated with the anti-SjPrx-4 mouse serum (1:200 in PBS) at 4 °C overnight in a humidified chamber. Non-specific antigen-antibody reaction was blocked with Histofine Mouse Stain Kit (Nichirei Biosciences, Tokyo, Japan) whereas Histofine Simplestain AP (Nichirei Biosciences) was used as the secondary antibody, both according to the manufacturer's instructions. Labeling was visualized with a Fast Red chromogen (Fast Red II Substrate Kit; Nichirei Biosciences), and the sections were counterstained with Meyer's hematoxylin. The sections incubated with normal mouse serum instead of the anti-SjPrx-4 mouse serum were used as negative controls.

#### ELISA with recombinant antigens

ELISAs were performed with rSjPrx-4, rSjTPx-1 and combination of rSjPrx-4 and rSjTPx-1 (mixed in equivalent molar) using a panel of 45 negative control samples from non-endemic USA, 41 schistosomiasis-negative and 30 schistosomiasis-positive samples confirmed through Kato-Katz from endemic municipalities in the Philippines. Sera from other parasitic infections, i.e. with *P. westermani*, *E. histolytica*, *O. viverrini*, *T. trichiura*, were used to check the cross-reactivity with the combination of rSjPrx-4/rSjTPx-1. Two hundred ng/well of the recombinant antigens or *S. japonicum* soluble egg antigen (SEA) diluted with carbonate/bicarbonate buffer (pH 9.6) were coated in the microplate plates (ThermoFisher Scientific, Roskilde, Denmark) overnight at 4 °C. After 3 washes with PBS containing 0.05% Tween 20 (PBS-T) (Wako, Osaka, Japan), blocking was performed with 120  $\mu$ l of 3% skimmed milk in PBS-T at RT for 5 min. Then, 100  $\mu$ l of serum samples at a dilution of 1:400 in blocking buffer was added to each well in triplicate and incubated for 1 h at 37 °C. After washing, the plates were incubated with 100  $\mu$ l of HRP-conjugated anti-human IgG (1:20,000) (Proteintech, Manchester, UK) as the secondary antibody at 37 °C for 1 h. Subsequently, the plates were washed 3 times with T-PBS and incubated with 100  $\mu$ l of HRP substrate, TMB (KPL, Gaithersburg, MD, USA) for 10 min. Fifty microliters of 1 M phosphoric acid was used to stop the reaction. The absorbance was measured at 450 nm using a microplate reader (MTP-500; Corona

Electric, Tokyo, Japan). The cut-off OD values were calculated from the values of 45 non-endemic controls as mean + 3SD.

### Statistical analysis

Data from qRT-PCR were represented as the mean  $\pm$  standard error (SE). Student's t-test was used to analyze data, with a *P*-value less than 0.05 considered significant. The online software MedCalc ([https://www.medcalc.org/calc/diagnostic\\_test.php](https://www.medcalc.org/calc/diagnostic_test.php)) was used to compute the sensitivity and specificity of the ELISA tests.

## Results

### Cloning, expression and purification of rSjPrx-4

The full-length coding sequence of SjPrx-4 was obtained by RT-PCR from the cDNA libraries of eggs and 8-week-old adult worms (Fig. 1a). The SjPrx-4 gene was 585 bp in size which was identical to the one described in GeneDB and coded a protein of 194 amino acids with a predicted molecular weight of 21.7 kDa and an isoelectric point of 7.6. The deduced polypeptide sequence of SjPrx-4 was 93.3% identical to its homologous gene SmPrx-4 in *S. mansoni* (GenBank: XP\_018651610.1) and 59.3–75.0% identity with SjTPx-3 (GenBank: BAD90103.1), SjTPx-2 (GenBank: BAD90102.1) and SjTPx-1 (GenBank: BAD01572.1) (Additional file 1: Table S1, Additional file 2: Figure S1). SjPrx-4 contained 2 cysteine conserved sequences with the peroxidatic cysteine residue at the N-terminal FYPADFTFVCPTE and the resolving cysteine residue at C-terminal GEVCPA as active sites (Additional file 1: Table S1, Additional file 2: Figure S1). The coding sequence of SjPrx-4 was successfully expressed as a GST fusion protein in *E. coli* Rosetta (DE3) (Fig. 1b, Lane 2). SDS-PAGE showed rSjPrx-4 with an expected size of 21.7 kDa after removing the GST tag (Fig. 1b, Lane 3).

### Transcription levels of SjPrx-4 gene at different developmental stages of *S. japonicum*


RT-qPCR was performed to examine the transcription levels of SjPrx-4 in eggs, sporocysts and 8-week-old male and female adult worms. SjPrx-4 transcript was found in all stages examined (Fig. 2a). SjPrx-4 transcription expression was highest level in males, followed by eggs and females (Fig. 2b). On the other hand, sporocysts showed the lowest transcription level of SjPrx-4 (Fig. 2b).

### Antioxidant activity of rSjPrx-4

The antioxidant activity of rSjPrx-4 was measured as the capacity in protection of supercoiled DNA from nicking by using the MFO system. The MFO system generated hydroxyl radicals that changed Bluescript supercoiled plasmid DNA to nicked form as confirmed by its increased molecular weight in the agarose gel (Fig. 3, Lane 4). However, the presence of rSjPrx-4 at concentrations of above 1  $\mu$ g/ml prevented the nicking of supercoiled plasmid DNA (Fig. 3, Lanes 7, 8). In addition, presence of heat-inactivated rSjPrx-4 did not prevent the nicking of supercoiled plasmid DNA (Fig. 3, lane 9, 10). These results confirmed the antioxidant activity of rSjPrx-4.

### Western blot analysis of anti-rSjPrx-4 against egg and adult worm lysates

Anti-rSjPrx-4 mouse serum was used to detect native SjPrx-4 in eggs and adult worms. The results showed the presence of a band in egg and adult worm lysates at approximately 21.7 kDa as the expected size of the protein (Fig. 4, Lanes 3, 5). The normal mouse serum showed no specific bands with the protein extract of egg and adult worms (Fig. 4, Lanes 2, 4).



### Immunolocalization of SjPrx-4 in *S. japonicum* eggs and adult worms

Immunohistochemical staining was conducted to observe the localization of SjPrx-4 in the eggs deposited in the infected liver sections and schistosome adult worms, using the mouse anti-rSjPrx-4 serum. Sections treated with the anti-rSjPrx-4 serum showed that SjPrx-4 was mainly localized inside the egg and in the parenchyma as well as on the tegument of adult male worms (Fig. 5a, c), as opposed to the sections incubated with normal mouse serum, which showed no positive reactions (Fig. 5b, d).

### ELISA with rSjPrx-4, rSjTPx-1 and a combination of SjPrx-4/SjTPx-1

SEA detected 27 out of 30 schistosomiasis-confirmed sera (90.0%) as positive and 29 out of 41 non-endemic sera (70.7%) as negative (Fig. 6, Additional file 3: Table S2). Individually, the SjPrx-4-ELISA showed 83.3% sensitivity and 87.6% specificity whereas the SjTPx-1-ELISA had 83.3% sensitivity and 97.6% specificity (Fig. 6, Additional file 3: Table S2). However, when rSjTPx-1 and rSjPrx-4 were combined as ELISA antigens, the sensitivity and specificity of the test improved to 90.0% and 92.7%, respectively (Fig. 6, Additional file 3: Table S2). Moreover, the results of combined SjTPx-1/SjPrx-4 ELISA analyzed by receiver-operating characteristic curves showed the best sensitivity of 97.6% and specificity of 90.0% (Additional file 4: Figure S2).


Human serum samples collected from the patients infected with *T. trichiura* and *E. histolytica* had no


cross-reaction in the ELISA with a combination of rSjPrx-4 and rSjTPx-1. However, 1 out of 6 *O. viverrini*-infected and 1 out of 10 *P. westermani*-infected human sera showed minimum cross-reaction against the combined antigen with OD values slightly higher than the cut-off value.

### Discussion


Molecular cloning and characterization of 2-Cys peroxiredoxins such as SjTPx-1, SjTPx-2 and SjTPx-3 from *S. japonicum* [19, 21, 25] as well as SmTPx-1 [33], SmTPx-2 and SmTPx-3 [34] from *S. mansoni* have been reported in previous research. A fourth type, SmPrx-4, was first identified by Protasio et al. [22] in 2012 when they systematically improved the *S. mansoni* draft genome using a combination of Sanger and Next Generation sequencing. Moreover, the Prx-4 homologue was also found in *S. mekongi* when egg proteins were analyzed using proteomic strategies [35]. In the present study, a novel Prx-4 homologue from *S. japonicum* (SjPrx-4) was successfully cloned and characterized.

We have obtained the SjPrx-4 complete sequence comprised of 585 bp that encoded 194 amino acids without a signal peptide and with a predicted molecular weight of 21.7 kDa. Bioinformatics analysis showed that SjPrx-4 was most closely related to SmPrx-4 exhibiting two conserved cysteines, the peroxidatic Cys and the resolving Cys. These cysteines of the 2-Cys Prx family are known to be essential in Cys-dependent thioredoxin peroxidase activity [36]. Moreover, the amino acid sequence of





**Fig. 3** Antioxidant activity of SjPrx-4. Lane 1: pBluescript plasmid DNA; Lane 2: pBluescript plasmid DNA+ FeCl<sub>3</sub>; Lane 3: pBluescript plasmid DNA+DTT; Lane 4: pBluescript plasmid DNA + FeCl<sub>3</sub> + DTT; Lanes 5–8: pBluescript plasmid DNA + FeCl<sub>3</sub> + DTT and 10 ng, 100 ng, 1 μg, and 5 μg/ml of rSjPrx-4, respectively; Lanes 9, 10: pBluescript plasmid DNA + FeCl<sub>3</sub> + DTT and 1 μg, and 5 μg/ml of denatured rSjPrx-4, respectively. The nicked form (NF) and supercoiled form (SF) are indicated on the right




**Fig. 4** Western blot analysis of *S. japonicum* egg and adult worm lysates with anti-SjPrx-4 mouse sera. Lane 1: protein marker; Lane 2: *S. japonicum* egg lysate incubated with normal mouse sera; Lane 3: *S. japonicum* egg lysate incubated with anti-SjPrx-4 mouse sera; Lane 4: adult worm lysate incubated with normal mouse sera; Lane 5: adult worm lysate incubated with anti-SjPrx-4 mouse sera

SjPrx-4, SjTPx-1 and SjTPx-2 contains the GVL motif, which was reported to be essential to the catalase-like activity of Prx-1 as seen in the green spotted puffer fish *Tetradon nigroviridis* [37]. The catalase-like activity might explain how some parasitic helminths such as

*Fasciola hepatica* and *S. mansoni* are deficient in catalase [38] and have relatively low levels of glutathione peroxidase [39], but possess highly expressed Prxs [40, 41]. SjPrx-4 as well as SjTPx-1 and SjTPx-2 may be enzymes having dual-functions by acting as Cys-dependent thioredoxin peroxidase and having a catalase-like activity at the same time. The catalase-like activity of SjPrx-4, SjTPx-1 and SjTPx-2 should however be confirmed with further studies.

The results of antioxidant activity showed Prx proteins from other helminths had similar activity in protecting supercoiled plasmid DNA in a MFO assay [34, 42]. In this study, rSjPrx-4 also showed antioxidant activity in protection of pBluescript from supercoiled DNA form to nicked form in MFO system. This finding suggested that SjPrx-4 might help the parasite to neutralize the ROS generated from the host immune system as well as from the internal metabolism of worms themselves.

The RT-qPCR results showed the presence of SjPrx-4 in all the stages of *S. japonicum* examined here. The finding that the expression of the SjPrx-4 transcript in male worms was higher than that of female worms may imply the fact that the latter require less SjPrx-4 expression since they are enclosed in the gynecophoral canal of the male worms. The present study showed that expression profile of SjPrx-4 was different from the three other TPxs. In contrast, RT-qPCR results of SjTPx-1, SjTPx-2



**Fig. 5** Immunolocalization of SjPrx-4 in eggs and adult worms of *S. japonicum*. The red arrows indicate the existence of SjPrx-4 expression. **a** The liver section from *S. japonicum*-infected mice reacted with anti-SjPrx-4 mouse serum shows immunopositivity in the space between the miracidium and the egg-shell. **b** There is no reaction in the liver section from *S. japonicum*-infected mice reacted with normal mouse serum. **c** Immunopositive reactions of SjPrx-4 were observed in the tegument (te) and in the parenchyma (p) of *S. japonicum* adult worms (♀: female and ♂: male). **d** The section from *S. japonicum* adult worms reacted with normal mouse serum did not reveal any immunopositive reactions. Scale-bars: 50 µm

and SjTPx-3 showed that their mRNA in adult female worms were more abundant than in adult male worms. It was suggested that adult female worms may express the three other TPxs at a higher level than adult male worms [21, 25]. The result of RT-qPCR of SjPrx-4 was also supported by the immunohistochemistry experiments in the present study, showing that SjPrx-4 was mainly localized in the tegument of adult male worms as compared to the adult female worms. The more exposed male worms in turn express high levels of this enzyme to protect themselves from oxidative stress posed by the host immune system.


Immunolocalization of SjPrx-4 revealed that the protein is distributed inside the egg-shell, in the parenchyma and in the tegument of the adult worm. These results are different from the other reported *S. japonicum* Prxs: SjTPx-1 is a tegumental and circulating antigen; SjTPx-2 exists in the sub-tegument, parenchyma, gut epithelia and vitelline gland; and SjTPx-3 is a tegumental antigen having a mitochondrial signal peptide [19, 21]. It was

suggested that the antioxidant functions of TPx-2 are associated with intracellular redox signaling [19] whereas those of SjTPx-3 are associated with the mitochondria [19, 21, 43]. The Prx family may have different roles at different locations and compensate with each other to protect the parasite from oxidative stresses. SjPrx-4 may have a chance to be exposed to the host's immune system during infection, and thus may be a potential antigen for serodiagnosis in detecting the parasite infection.

The western blot results were consistent with those of immunolocalization and RT-qPCR in the egg and adult worm stages. The appearance of a single band showed that anti-SjPrx-4 can specifically recognize the native SjPrx-4 of *S. japonicum*.

The rSjPrx-4 and rSjTPx-1 used as a single antigen in the ELISA showed a positive rate of 83.3% in comparison with the 90% sensitivity of SEA. However, the specificity of rSjPrx-4 and SjTPx-1 was higher (at 87.8% and 97.6%, respectively) than that of SEA at 70.7%. Although both rSjPrx-4 and rSjTPx-1 were shown to be potential





diagnostic antigens, their sensitivity and specificity need to be further improved to meet the current needs of schistosomiasis diagnosis. A way to improve the diagnostic capabilities of ELISA in parasitic diseases is combining multiple antigens in a cocktail format. Some of the patient sera have high OD values with SjtPx-4 but not with SjtPx-1 and *vice versa* (Additional file 5: Figure S3). Hence, two antigens might complement each other to have a better synergistic diagnostic potential. The combination of SjtPx-1 and SjtPx-4 antigens has improved the sensitivity of the ELISA to 90.0%. There have already been some reports on the application of multi-epitope recombinant antigens (chimeric proteins) to improve the sensitivity of serodiagnostic tests for schistosomiasis in experimental animals, goats and water buffaloes [44–46]. Therefore, it appears possible to construct multi-epitope recombinant antigens of SjtPx-1 and SjtPx-4 to enhance the accuracy and reliability of the diagnostic tools in detecting human schistosomiasis.

## Conclusions

Our study suggests that SjtPx-4 may play a role as an antioxidant of *S. japonicum* to deal with oxidative stress and that rSjtPx-4 could be a potential antigen for use

in ELISA-based diagnosis of *S. japonicum* infection in humans. The combination of SjtPx-4 and SjtPx-1 as sensitive complementary diagnostic antigens suggests its use in the development of fusion antigens that could improve the sensitivity of ELISA assays in detecting human *S. japonicum* infection.

## Supplementary information

**Supplementary information** accompanies this paper at <https://doi.org/10.1186/s13071-020-04313-w>.

**Additional file 1: Table S1.** Identity of the SjtPx-4 amino acid sequence with SmPrx-4, SjtPx-1, SjtPx-2 and SjtPx-3.

**Additional file 2: Figure S1.** Amino acid sequence alignment of SjtPx-4 with SmPrx-4, SjtPx-1, SjtPx-2 and SjtPx-3.

**Additional file 3: Table S2.** Statistical analysis of ELISA results of rSjtPx-1, rSjtPx-4 and a combination of rSjtPx-4/rSjtPx-1.

**Additional file 4: Figure S2.** Receiver operating characteristic (ROC) curves for ELISA of SEA, SjtPx-4, SjtPx-1 and combination of SjtPx-1/SjtPx-4 against sera from endemic negative ( $n = 41$ ) and schistosomiasis-positive by Kato-Katz ( $n = 30$ ).

**Additional file 5: Figure S3.** Kinetics of 30 *S. japonicum* stool-positive sera response to SjtPx-1 and SjtPx-4.

## Abbreviations

Cq: threshold cycle; ELISA: enzyme-linked immunosorbent assay; GST: glutathione S-transferase; HRP: horseradish peroxidase; ICR: Institute of Cancer

Research; IPTG: isopropyl- $\beta$ -D-thiogalactoside; MFO: mixed-function oxidation; Prx: peroxiredoxin; PVDF: polyvinylidene difluoride; qRT-PCR: quantitative real time polymerase chain reaction; SE: standard error; SDS-PAGE: sodium dodecyl sulfate polyacrylamide gel electrophoresis; TBS-T: tris-buffered saline containing 0.05% Tween 20; TPI: triose-phosphate isomerase; TPx: thioredoxin peroxidase; PBS-T: phosphate-buffered saline containing 0.05% Tween 20.

#### Acknowledgements

We are grateful to Dr Haruhiko Maruyama, Miyazaki University Graduate School of Medicine, Miyazaki, Japan, for the *P. westermani* positive samples, and Dr. Isao Nagano, Department of Parasitology, Gifu University Graduate School of Medicine, Gifu, Japan, for providing the *O. viverrini*-positive samples.

#### Authors' contributions

MADT, JMMA, YG and SK designed and conceived the study. MADT carried out the experiments, analyzed the data and drafted the manuscript. AMCM, TTN, KJM, JMMA and SK participated in revising the manuscript. MADT, AMCM, TTN and LH carried out the animal experiments. YC and MK contributed to the animal experiments. MADT and SN performed the immunohistochemistry analysis. KJM, JMMA, YG and SK collected human serum samples. All authors read and approved the final manuscript.

#### Funding

This study was supported partly by a grant from Japan Society for the Promotion of Science to SK (19KK0173).

#### Availability of data and materials

Data supporting the conclusions of this article are included within the article and its additional files.

#### Ethics approval and consent to participate

Patient sera in the Philippines were carried out according to the ethical clearance sought from the University of the Philippines, Manila (clearance no: UPM REB 2011-098) and Obihiro University of Agriculture and Veterinary Medicine (Approval no: 2010-03-02). All animal experiments were carried out in accordance with the guidelines for the Care and Use of Laboratory Animals established by Obihiro University of Agriculture and Veterinary Medicine, and Dokkyo Medical University. This study was approved by the Committee for Animal Experimentation of Obihiro University of Agriculture and Veterinary Medicine, and Dokkyo Medical University (Approval no. 29-52 and 0006).

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare that they have no competing interests.

#### Author details

<sup>1</sup> National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Hokkaido, Japan. <sup>2</sup> The United Graduate School of Veterinary Sciences, Gifu University, Gifu, Japan. <sup>3</sup> Department of Immunology and Microbiology, Pasteur Institute in Ho Chi Minh City, Ho Chi Minh City, Vietnam. <sup>4</sup> Department of Parasitology, College of Public Health, University of the Philippines Manila, Manila, Philippines. <sup>5</sup> Department of Biology, School of Science and Engineering, Ateneo de Manila University, Manila, Philippines. <sup>6</sup> Department of Immunopathology and Microbiology, College of Veterinary Medicine and Biomedical Sciences, Cavite State University, Cavite, Philippines. <sup>7</sup> Laboratory of Veterinary Pathology, Obihiro University of Agriculture and Veterinary Medicine, Hokkaido, Japan. <sup>8</sup> Department of Tropical Medicine and Parasitology, Dokkyo Medical University, Tochigi, Japan. <sup>9</sup> Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo, Japan.

Received: 6 May 2020 Accepted: 24 August 2020

Published online: 01 September 2020

#### References

- Bergquist R, Johansen MV, Utzinger J. Diagnostic dilemmas in helminthology: what tools to use and when? *Trends Parasitol.* 2009;25:151–6.

- WHO. Prevention and control of schistosomiasis and soil-transmitted helminthiasis. Geneva: World Health Organization; 2002.
- Lier T, Simonsen GS, Wang T, Lu D, Haukland HH, Vennervald BJ, et al. Low sensitivity of the formol-ethyl acetate sedimentation concentration technique in low-intensity *Schistosoma japonicum* infections. *PLoS Negl Trop Dis.* 2009;3:e386.
- Lin DD, Liu JX, Liu YM, Hu F, Zhang YY, Xu JM, et al. Routine Kato-Katz technique underestimates the prevalence of *Schistosoma japonicum*: a case study in an endemic area of the People's Republic of China. *Parasitol Int.* 2008;57:281–6.
- Zhang YY, Luo JP, Liu YM, Wang QZ, Chen JH, Xu MX, et al. Evaluation of Kato-Katz examination method in three areas with low-level endemicity of schistosomiasis japonica in China: a Bayesian modeling approach. *Acta Trop.* 2009;112:16–22.
- Liu S, Zhou X, Piao X, Hou N, Shen Y, Zou Y, et al. Saposin-like proteins, a multigene family of *Schistosoma* species, are biomarkers for the immunodiagnosis of schistosomiasis japonica. *J Infect Dis.* 2016;214:1225–34.
- Xu X, Zhang Y, Lin D, Zhang J, Xu J, Liu YM, et al. Serodiagnosis of *Schistosoma japonicum* infection: genome-wide identification of a protein marker, and assessment of its diagnostic validity in a field study in China. *Lancet Infect Dis.* 2014;14:489–97.
- Cai P, Weerakoon KG, Mu Y, Olveda DU, Piao X, Liu S, et al. A parallel comparison of antigen candidates for development of an optimized serological diagnosis of schistosomiasis japonica in the Philippines. *EBioMedicine.* 2017;24:237–46.
- Houghton RL, Benson DR, Reynolds LD, McNeill PD, Sleath PR, Lodes MJ, et al. A multi-epitope synthetic peptide and recombinant protein for the detection of antibodies to *Trypanosoma cruzi* in radioimmuno-precipitation-confirmed and consensus-positive sera. *J Infect Dis.* 1999;179:1226–34.
- Li S, Shin JG, Cho PY, Kim TI, Hong ST, Hong SJ. Multiple recombinant antigens of *Clonorchis sinensis* for serodiagnosis of human clonorchiasis. *Parasitol Res.* 2011;108:1295–302.
- Kim S, Ahn HJ, Kim TS, Nam HW. ELISA detection of vivax malaria with recombinant multiple stage-specific antigens and its application to survey of residents in endemic areas. *Korean J Parasitol.* 2003;41:203–7.
- Moendeg KJ, Angeles JM, Goto Y, Leonardo LR, Kirinoki M, Villacorte EA, et al. Development and optimization of cocktail-ELISA for a unified surveillance of zoonotic schistosomiasis in multiple host species. *Parasitol Res.* 2015;114:1225–8.
- Vieira P, Miranda HP, Cerqueira M, Delgado Mde L, Coelho H, Antunes D, et al. Latent schistosomiasis in Portuguese soldiers. *Mil Med.* 2007;172:144–6.
- Harris AR, Russell RJ, Charters AD. A review of schistosomiasis in immigrants in western Australia, demonstrating the unusual longevity of *Schistosoma mansoni*. *Trans R Soc Trop Med Hyg.* 1984;78:385–8.
- Loverde PT. Do antioxidants play a role in schistosome host-parasite interactions? *Parasitol Today.* 1998;14:284–9.
- Chae HZ, Kim IH, Kim K, Rhee SG. Cloning, sequencing, and mutation of thiol-specific antioxidant gene of *Saccharomyces cerevisiae*. *J Biol Chem.* 1993;268:16815–21.
- Kang SW, Baines IC, Rhee SG. Characterization of a mammalian peroxiredoxin that contains one conserved cysteine. *J Biol Chem.* 1998;273:6303–11.
- Dzik JM. Molecules released by helminth parasites involved in host colonization. *Acta Biochim Pol.* 2006;53:33–64.
- Kumagai T, Osada Y, Kanazawa T. 2-Cys peroxiredoxins from *Schistosoma japonicum*: the expression profile and localization in the life cycle. *Mol Biochem Parasitol.* 2006;149:135–43.
- Macalanda AMC, Angeles JMM, Moendeg KJ, Dang AT, Higuchi L, Inoue N, et al. Evaluation of *Schistosoma japonicum* thioredoxin peroxidase-1 as a potential circulating antigen target for the diagnosis of Asian schistosomiasis. *J Vet Med Sci.* 2018;80:156–63.
- Han Y, Zhao B, Zhang M, Hong Y, Han H, Cao X, et al. Biochemical properties and vaccine effect of recombinant TPx-3 from *Schistosoma japonicum*. *Parasitol Res.* 2017;116:1361–72.
- Protasio AV, Tsai JJ, Babbage A, Nichol S, Hunt M, Aslett MA, et al. A systematically improved high quality genome and transcriptome of the human blood fluke *Schistosoma mansoni*. *PLoS Negl Trop Dis.* 2012;6:e1455.

23. Zhang W, Rogniaux H, Huang W, Chauvin A, Moreau E. Analysis of thioredoxin peroxidase as a promising antigen for diagnosis of *Fasciola gigantica* infection: a preliminary study. *Parasitol Int.* 2011;60:206–8.
24. Hakimi H, Goto Y, Suganuma K, Angeles JM, Kawai S, Inoue N, et al. Development of monoclonal antibodies against *Plasmodium falciparum* thioredoxin peroxidase 1 and its possible application for malaria diagnosis. *Exp Parasitol.* 2015;154:62–6.
25. Hong Y, Han Y, Fu Z, Han H, Qiu C, Zhang M, et al. Characterization and expression of the *Schistosoma japonicum* thioredoxin peroxidase-2 gene. *J Parasitol.* 2013;99:68–76.
26. Angeles JM, Goto Y, Kirinoki M, Leonardo L, Tongol-Rivera P, Villacorte E, et al. Human antibody response to thioredoxin peroxidase-1 and tandem repeat proteins as immunodiagnostic antigen candidates for *Schistosoma japonicum* infection. *Am J Trop Med Hyg.* 2011;85:674–9.
27. Angeles JM, Goto Y, Kirinoki M, Asada M, Leonardo LR, Rivera PT, et al. Utilization of ELISA using thioredoxin peroxidase-1 and tandem repeat proteins for diagnosis of *Schistosoma japonicum* infection among water buffaloes. *PLoS Negl Trop Dis.* 2012;6:e1800.
28. Angeles JMM, Goto Y, Kirinoki M, Leonardo LR, Moendeg KJ, Ybanez AP, et al. Detection of canine *Schistosoma japonicum* infection using recombinant thioredoxin peroxidase-1 and tandem repeat proteins. *J Vet Med Sci.* 2019;10:1413–8.
29. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-</sup>(Delta Delta C(T)) method. *Methods.* 2001;25:402–8.
30. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc.* 2008;3:1101–8.
31. Sauri H, Butterfield L, Kim A, Shau H. Antioxidant function of recombinant human natural killer enhancing factor. *Biochem Biophys Res Commun.* 1995;208:964–9.
32. Ramos-Vara JA, Kiupel M, Baszler T, Bliven L, Brodersen B, Chelack B, et al. Suggested guidelines for immunohistochemical techniques in veterinary diagnostic laboratories. *J Vet Diagn Invest.* 2008;20:393–413.
33. Kwatia MA, Botkin DJ, Williams DL. Molecular and enzymatic characterization of *Schistosoma mansoni* thioredoxin peroxidase. *J Parasitol.* 2000;86:908–15.
34. Sayed AA, Williams DL. Biochemical characterization of 2-Cys peroxiredoxins from *Schistosoma mansoni*. *J Biol Chem.* 2004;279:26159–66.
35. Thiangtrongjit T, Adisakwattana P, Limpanont Y, Dekumyoy P, Nuanmtanong S, Chusongsang P, et al. Proteomic and immunomic analysis of *Schistosoma mekongi* egg proteins. *Exp Parasitol.* 2018;191:88–96.
36. Park J, Lee S, Lee S, Kang SW. 2-cys peroxiredoxins: emerging hubs determining redox dependency of mammalian signaling networks. *Int J Cell Biol.* 2014;2014:715867.
37. Sun CC, Dong WR, Shao T, Li JY, Zhao J, Nie L, et al. Peroxiredoxin 1 (Prx1) is a dual-function enzyme by possessing Cys-independent catalase-like activity. *Biochem J.* 2017;474:1373–94.
38. Mkoji GM, Smith JM, Prichard RK. Antioxidant systems in *Schistosoma mansoni*: correlation between hydrogen susceptibility to oxidant killing and the levels of scavengers of hydrogen peroxide and oxygen free radicals. *Int J Parasitol.* 1988;18:661–6.
39. Maiorino M, Roche C, Kiess M, Koenig K, Gawlik D, Matthes M, et al. A selenium-containing phospholipid-hydroperoxide glutathione peroxidase in *Schistosoma mansoni*. *Eur J Biochem.* 1996;238:838–44.
40. Robinson MW, Hutchinson AT, Dalton JP, Donnelly S. Peroxiredoxin: a central player in immune modulation. *Parasite Immunol.* 2010;32:305–13.
41. Mei H, LoVerde PT. *Schistosoma mansoni*: the developmental regulation and immunolocalization of antioxidant enzymes. *Exp Parasitol.* 1997;86:69–78.
42. Suttiprapa S, Loukas A, Laha T, Wongkham S, Kaewkes S, Gaze S, et al. Characterization of the antioxidant enzyme, thioredoxin peroxidase, from the carcinogenic human liver fluke. *Opisthorchis viverrini*. *Mol Biochem Parasitol.* 2008;160:116–22.
43. Sayed AA, Cook SK, Williams DL. Redox balance mechanisms in *Schistosoma mansoni* rely on peroxiredoxins and albumin and implicate peroxiredoxins as novel drug targets. *J Biol Chem.* 2006;281:17001–10.
44. Jin YM, Lu K, Zhou WF, Fu ZQ, Liu JM, Shi YJ, et al. Comparison of recombinant proteins from *Schistosoma japonicum* for schistosomiasis diagnosis. *Clin Vaccine Immunol.* 2010;17:476–80.
45. Lv C, Hong Y, Fu Z, Lu K, Cao X, Wang T, et al. Evaluation of recombinant multi-epitope proteins for diagnosis of goat schistosomiasis by enzyme-linked immunosorbent assay. *Parasit Vectors.* 2016;9:135.
46. Lv C, Fu Z, Lu K, Yue R, Wang T, Cao X, et al. A perspective for improving the sensitivity of detection: the application of multi-epitope recombinant antigen in serological analysis of buffalo schistosomiasis. *Acta Trop.* 2018;183:14–8.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.