

RESEARCH

Open Access

Immunoregulation of bovine macrophages by factors in the salivary glands of *Rhipicephalus microplus*

Danett K Brake* and Adalberto A Pérez de León

Abstract

Background: Alternative strategies are required to control the southern cattle tick, *Rhipicephalus microplus*, due to evolving resistance to commercially available acaricides. This invasive ectoparasite is a vector of economically important diseases of cattle such as bovine babesiosis and anaplasmosis. An understanding of the biological intricacies underlying vector-host-pathogen interactions is required to innovate sustainable tick management strategies that can ultimately mitigate the impact of animal and zoonotic tick-borne diseases. Tick saliva contains molecules evolved to impair host innate and adaptive immune responses, which facilitates blood feeding and pathogen transmission. Antigen presenting cells are central to the development of robust T cell responses including Th1 and Th2 determination. In this study we examined changes in co-stimulatory molecule expression and cytokine response of bovine macrophages exposed to salivary gland extracts (SGE) obtained from 2-3 day fed, pathogen-free adult *R. microplus*.

Methods: Peripheral blood-derived macrophages were treated for 1 hr with 1, 5, or 10 µg/mL of SGE followed by 1, 6, 24 hr of 1 µg/mL of lipopolysaccharide (LPS). Real-time PCR and cytokine ELISA were used to measure changes in co-stimulatory molecule expression and cytokine response.

Results: Changes were observed in co-stimulatory molecule expression of bovine macrophages in response to *R. microplus* SGE exposure. After 6 hrs, CD86, but not CD80, was preferentially up-regulated on bovine macrophages when treated with 1 µg/ml SGE and then LPS, but not SGE alone. At 24 hrs CD80, CD86, and CD69 expression was increased with LPS, but was inhibited by the addition of SGE. SGE also inhibited LPS induced upregulation of TNFα, IFNγ and IL-12 cytokines, but did not alter IL-4 or CD40 mRNA expression.

Conclusions: Molecules from the salivary glands of adult *R. microplus* showed bimodal concentration-, and time-dependent effects on differential up-regulation of CD86 in bovine macrophages activated by the TLR4-ligand, LPS. Up regulation of proinflammatory cytokines and IL-12, a Th1 promoting cytokine, were inhibited in a dose-dependent manner. The co-stimulatory molecules CD80, as well as the cell activation marker, CD69, were also suppressed in macrophages exposed to SGE. Continued investigation of the immunomodulatory factors will provide the knowledge base to research and develop therapeutic or prophylactic interventions targeting *R. microplus*-cattle interactions at the blood-feeding interface.

Background

Ticks are external parasitic organisms that have to overcome host defence mechanisms to obtain blood for their survival. They also serve as vectors of pathogens causing important diseases in animals and humans [1]. As a

result, complex tick-pathogen-host interactions have developed through evolutionary time. The southern cattle tick, *Rhipicephalus (Boophilus) microplus*, is one of the most economically important parasites of livestock. Additionally, this invasive tick species is also a significant vector of *Babesia bigemina* and *B. bovis* that cause bovine babesiosis, which can be deadly to cattle [2]. Alternative strategies are required to control *R. microplus* as populations across the globe continue to evolve

* Correspondence: danett.brake@ars.usda.gov
USDA-ARS Knippling-Bushland U.S. Livestock Insects Research Laboratory, 2700
Fredericksberg Rd, Kerrville, TX 78028 USA

resistance to commercially available acaricides [3,4]. An understanding of the biological intricacies underlying vector-host-pathogen interactions, including those involving the host immune system, is required to innovate sustainable tick management strategies that can ultimately mitigate the impact of animal and zoonotic tick-borne diseases.

Tissue injury at the tick feeding site activates the different arms of the host immune system. While feeding, ticks secrete bioactive salivary factors to modulate humoral and cellular components of the innate and acquired immune responses to improve reproductive fitness [5,6]. This immunomodulation by salivary factors has been shown to affect the activity of antigen presenting cells (APC's), lymphocytes and other cells, and to inhibit complement activation [7-11]. These effects on the host immune system may also enhance the transmission of tick-borne pathogens [5].

Many types of APC's including macrophages, different subtypes of dendritic cells (DC), and natural killer (NK) cells reside in the skin and play a crucial role in inducing protective T cell responses. Langerhans cells (LC), a type of DC found in the skin, from guinea pig epidermis have been shown to acquire antigens from tick salivary glands, migrate from infestation sites and present them at local lymph nodes [12,13]. Tick-sensitized animals have greater numbers of LC at sites of tick infestation [14].

How tick saliva alters antigen presenting cell function is not well understood. Inflammatory signals can effect DC homeostasis, activation and differentiation [15]. Several studies describe the effects of saliva, or salivary gland extracts (SGE) on proinflammatory cytokine expression in murine *in vitro* models. Studies with *Dermacentor andersoni*, *Ixodes pacificus*, *I. ricinus*, and *R. sanguineus* have shown a tick-induced shift away from Th1 cytokines such as tumor necrosis factor α (TNF α), interferon γ (IFN γ), and interleukin 1 β (IL-1 β), to promoting up-regulation of interleukin-10 (IL-10), and interleukin-4 (IL-4), which are consistent with Th2 polarization [11,16-21]. Specifically, a sphingomyelinase-like enzyme has been identified in *I. scapularis* that reduces antigen specific responses and promotes Th2 polarization [18,22].

Tick saliva may direct DC differentiation and function to drive naïve CD4 T cells towards Th2 differentiation [16,23]. Mice deficient in Langerhans cells, a subset of skin DCs, prevent the suppression of a Th1 response when exposed to *I. scapularis* ticks [24]. Salivary prostaglandin E₂ from *I. scapularis* can also suppress CD4 T cell proliferation by *in vitro* derived dendritic cells [25]. Altering the host immune response to a Th2 phenotype may benefit the transmission by *R. microplus* of pathogens like the apicomplexan protozoa causing bovine

babesiosis, which would be more successfully controlled by a Th1 response [26].

Another possible route to suppress APC's ability to induce a robust T cell response is by inhibiting cell maturation and activation. Upon activation, APCs up-regulate CD80 and CD86, which via binding to CD28 provide co-stimulatory signals for T cell activation [27]. CD80 and CD86 can modulate naïve T cells towards Th1 or Th2 pathways [28-30]. While CD80 preferentially favors Th1 type T cell differentiation, CD86 promotes IL-4 production and Th2 T cell differentiation [29,31,32]. CD86 can be differentially regulated by various cytokines including the Th2 cytokine, IL-4 [33]. Using a murine macrophage cell line, we demonstrated that molecules in SGE of *R. microplus* have a concentration-dependent effect on differential up-regulation of CD86 by the TLR4 ligand, LPS [34]. This CD86 up-regulation is at least partially dependent on the ERK1/2 pathway, and may serve to promote Th2 polarization of the immune response. Here, we investigated the effects of *R. microplus* salivary gland extract (SGE) on co-stimulatory molecule expression, macrophage activation, and cytokine expression in cultured bovine macrophages.

Methods

Isolation of Tick Salivary Glands

The *R. microplus* Deutch strain used as the source of ticks for this study originated from samples collected in Webb County, TX during an outbreak in 2001. The Deutch strain has been maintained by standard rearing practices at the USDA-ARS Cattle Fever Tick Research Laboratory at Moore Field, TX. The ticks were determined free of *Babesia bovis* and *Babesia bigemina* as described previously [35] and SGE was isolated as previously described [34]. Larvae were placed in patches and allowed to feed following protocols approved by the Institutional Animal Care and Use Committee of the USDA-ARS Knippling-Bushland Livestock Insects Research Laboratory. Adult ticks were fed for 2-3 days, cleaned with 70% ethanol and dissected. Salivary glands were sonicated at 55 kHz for three 20 second pulses on ice in Dulbecco's phosphate buffered saline (PBS) (Sigma, St. Louis, MO) and centrifuged at 14,000 \times g for 20 minutes at 4°C. The supernatant was collected and protein concentration was determined by Pierce BCA (bicinchoninic acid) Protein Assay (Thermo Scientific, Rockford, IL). The SGE was aliquoted and subsequently frozen at -70°C.

Isolation and culture of bovine macrophages

Whole blood was drawn from healthy Hereford cows following protocols approved by the Institutional Animal Care and Use Committee of the USDA-ARS Knippling-Bushland Livestock Insects Research Laboratory. Peripheral blood mononuclear cells (PBMC's) were isolated by

whole blood centrifugation using histopaque-1077 (Sigma-Aldrich, St. Louis, MO). PBMC's were seeded onto 24-well tissue culture treated plates in serum free RPMI 1640 (Gibco-Invitrogen, Carlsbad, CA). Monocytes were allowed to attach for 3 hrs at 37°C. Non-adherent cells were removed by serial washes with Dulbecco's phosphate-buffered saline (PBS) (Sigma, St. Louis, MO). Antibiotic-free RPMI media supplemented with 10% heat inactivated FBS was exchanged every 3-4 days. After 8-10 days, > 95% of cells had morphological characteristics consistent with a macrophage phenotype.

Real-time Quantitative PCR

Bovine *in vitro* differentiated macrophages were treated for 1 hr with 0, 1, 5 or 10 µg/mL of SGE followed by 1 µg/mL LPS or no additional treatment. After 1, 6 or 24 hrs of LPS treatment total RNA was extracted by spin column centrifugation using the RNAeasy Mini Kit (Qiagen, Valencia, CA). RNA concentration and absorbance at 260/280 nm was determined using a NanoDrop spectrophotometer (Thermo Scientific, Wilmington, DE) and RNA quality was analysed by non-denaturing agarose gel electrophoresis. cDNA synthesis and real-time PCR was performed as previously described [34]. For the amplification of specific mRNA, pre-inventoried 20× TaqMan MGB probe-primer sets for bovine TNF and GAPDH were purchased (Applied Biosystems, Foster City, CA). For other bovine genes, custom probe-primer sets were designed as shown on Table 1 (Sigma-Aldrich, St. Louis, MO). PCR was performed in a CFX96 Real-Time PCR Detection System (BioRad, Hercules, CA). Reactions were performed in duplicate. Relative mRNA expression was calculated by comparative C_t -method [36]. GAPDH was used as the endogenous control.

ELISA

Bovine *in vitro* differentiated macrophages were treated for 1 hr with 0, 1, 5 or 10 µg/mL of SGE followed by 1 µg/mL LPS for 24 hours. Cell culture supernatant was collected and quantities of TNFα and IFN-γ were assessed using bovine ELISA kits (R&D Systems,

Minneapolis, MN). Briefly, undiluted supernatant was incubated in the ELISA plate for 2 hrs, followed by washing and incubation with secondary reagents as supplied. Plates were read using a Spectramax Plus Microplate Spectrophotometer and Softmax Pro analysis software (Molecular Devices, Sunnyvale, CA).

Statistics

Results of 3-4 independent experiments are expressed as means ± SE. Significant differences between means were determined using unpaired Student's t-tests, or two-way analysis of variance (ANOVA) with $P < 0.05$ considered statistically significant.

Results

Changes in co-stimulatory molecule expression induced by SGE

Co-stimulatory molecule mRNA profiles of *in vitro* differentiated bovine macrophages were assayed by quantitative real-time PCR after 1, 6, or 24 hrs of treatment with or without 1, 5, or 10 µg/mL of SGE from adult ticks fed on cattle for 3 days, and 1 µg/mL of LPS. CD80, CD86, CD40 and CD69 were up-regulated in the presence of LPS (Figure 1). SGE alone did not alter message expression compared to untreated control at any concentration tested. 5 µg/mL SGE alone is represented in Figures 1 & 2. The 1 hr pre-treatment of SGE did reduce LPS induced up-regulation of CD80 and CD69 message in a dose dependent fashion ($P \leq 0.05$); however, CD40 expression was not significantly changed. Bovine macrophages treated with 1 µg/mL SGE and LPS showed increased CD86 expression at 6 hrs as compared to LPS alone, SGE alone or untreated cells. All concentrations of SGE treatments with LPS showed decreased CD86 expression after 24 hrs as compared to LPS alone.


Changes in cytokine expression following exposure to SGE

Cytokine mRNA expression levels were measured at 1, 6 and 24 hrs after LPS stimulation in the presence or

Table 1 Genes analysed by Taqman quantitative real-time PCR

Gene name	NCBI Accession	Sense	Anti-sense	Probe: FAM-TAMRA
CD86	NM_001038017	TAAGGCCGACAGCAGTTTCC	TCACCCCGTTATTAAGATGATAGC	TCCCAGCTCTGCTCCAGTCGGGT
CD80	Y09950	GCATTGTGATCCTGGCTCTG	CTGATCATTAACTCAGGAAGTC	TGTCGGACAGTGGCACCTACACCT
CD40	NM_001105611	GCATTGTGATCCTGGCTCTG	CTGATCATTAACTCAGGAAGTC	TGTCGGACAGTGGCACCTACACCT
CD69	NM_174014	ACCTTGGCCAAAACCTTTTGC	CAGCCCGATCCAGTGTTCAG	AACATGGTGCCACGCTTGCTGTCA
IFN-γ	NM_174086	GAAAGCGGAAGAGAAGTCAGAATC	CAAATATTGCAGGCAGGAGGAC	ACGTTGATGCTCTCCGGCCTCGAA
IL-12p40	NM_174356	TGTGACACTCCTGAAGAAGATGG	CCAAACTCTTTGACTTGGATGGTC	TGCCAGAGCCCAAGACCTCACTGC
IL-10	NM_174088	CTCTGTTGCCTGGTCTTCTCTG	TGGTTGGCAAGTGGATACAGC	CAGCCAGCCGAGATGCGAGCACCT
IL-4	NM_173921	AGCAAGACCTGTTCTGTGAATG	CAGCTTCAACACTTGGAGTATTTC	AGCCAAGACGAGCACAAGTACGCT

Gene names, NCBI gene accession, sense and anti-sense primer sequences (5' to 3') and Taqman probe sequence.




absence of varying SGE concentrations (Figure 2). No significant differences in IL-4 message were detected with SGE alone, LPS alone, and LPS in combination with SGE at any concentration tested. However, TNF, IFN- γ , and IL-12 mRNA expression was significantly decreased in LPS with SGE groups in a dose-dependent fashion when compared to LPS alone, SGE alone, or unstimulated control cells. Protein expression of TNF α and IFN- γ were measured in cell supernatant after 24 hrs of LPS stimulation (Figure 3). TNF α and IFN- γ were not detected in the control group, or the groups exposed to 1, 5, and 10 µg/mL SGE without LPS stimulation. A dose-dependent decrease of cytokine expression was observed with the addition of SGE when stimulated with LPS. Taken together, these data indicate *R. microplus* SGE contains factor(s) that suppress LPS-induced stimulation of macrophage activation, and cytokine responses.

Discussion

Cattle infested with *R. microplus* have been shown to reduce circulating bovine T and B cell numbers, and decreased cell responsiveness [37]. By comparison to innately resistant *B. indicus* cattle, less resistant *B.*


taurus breeds showed reduced numbers of basophils and eosinophils. Bovine microarray studies have shown that *R. microplus* differentially alters gene expression in susceptible *B. taurus* cattle as compared to *Bos indicus* breeds [38,39]. The expression of hemostatic proteins and adhesion molecules controlling coagulation and the recruitment of immune cells into sites of infestation, respectively, was also altered [40,41].

This study expands our investigations documenting how SGE from adult *R. microplus* affects the host's ability to mount a successful immune response by altering co-stimulatory and activation marker expression of bovine macrophages *in vitro*. Similar to our previous report using a mouse macrophage cell line [34], bovine macrophage CD40, which binds CD40L on T cells, was not significantly affected by exposure to SGE. At relatively low physiologic concentrations of SGE, as observed in our previous report, CD86 is up-regulated after 6 hours of LPS stimulation, but this effect was not seen at higher SGE concentrations. This finding is consistent with concentration-dependent bi-modal responses to differing levels of exposure to tick salivary proteins in the skin microenvironment and systemic responses. Low to moderate infestations with *R.*



microplus has been shown to induce an IgE response in cattle, whereas high infestations show increases in IgG antibody production [42]. CD86 expressed on human B cells can also promote IgE production when stimulated with IL-4 [43].

In contrast to results obtained with murine RAW 264.7 cells, *in vitro* differentiated bovine macrophages demonstrated a dramatic decrease in CD80 and CD86 mRNA expression by 24 hours of exposure to 5 and 10 µg/mL of SGE in the presence of LPS. These divergent



results highlight the differences in response between murine and bovine systems. Higher concentrations of saliva from adult *R. sanguineus* females fed for seven days can inhibit differentiation and maturation of murine bone-marrow-derived dendritic cells including CD80 and CD86 expression [23]. *I. ricinus* saliva pulsed dendritic cells were first reported to drive a Th2 response using 15 µg/mL saliva from females fed for 5.5 days [16,17]. In the presence of IL-1β, these DCs showed increased CD80 and CD86 expression and increased IL-4 production leading to Th2 priming of naïve CD4 T cells. Prostaglandin E₂ from *I. scapularis* saliva was shown to increase CD86 expression of LPS stimulated murine bone-marrow derived DC, while it inhibited IL-12p70 and TNFα protein expression in culture supernatants [25]. We did not observe changes in IL-4, but TNFα, IL-12p40, IFN-γ, CD80 and CD86 mRNA expression were decreased following exposure to *R. microplus* SGE at concentrations ≤ 10 µg/mL. These findings could be explained by differing immunoregulatory proteomes among species, varying concentration of salivary components tested, as well as changing compositions of salivary gland protein profiles during blood feeding [11,44-46].

These data support the hypothesis that in addition to altering APC function, the overall maturation and activation may be actively suppressed by molecules present in salivary glands of *R. microplus*. During the process of maturation macrophages and immature DCs change from phagocytic antigen processors to highly efficient activators of T cells. Up-regulation of CD80 and CD86 are markers of cell maturation as well as CD69. CD69 is expressed following activation in all bone marrow-derived cells [47]. The role of CD69 in the activation and regulation of the bovine immune response to tick salivary molecules is unknown. Our results document that CD69 expression is significantly reduced when SGE is present prior to LPS activation. Thus, tick salivary factors may actively prevent APC maturation needed for proper T cell activation. Additionally, CD69 is a complex co-stimulatory and immunoregulatory molecule as reviewed in [48]. CD69 may act as a proinflammatory receptor, promoting TNFα, nitric oxide secretion and IL-2 production, which increase T-cell proliferation. Knockout of CD69 in a murine asthma model showed increased eosinophil recruitment and enhanced Th2 cytokines in lung tissue [49]. Therefore CD69 may play a role in limiting Th2 responses. Further studies are needed to elucidate the role of CD69 in tick immunoregulation of host responses.

In adult *B. taurus*, ticks modulate host immune responses away from a Th1 profile by decreasing IFNγ, while promoting a Th2 phenotype via increased IL-4 production [18,19,23]. Moreover, this Th2 response

appears to facilitate pathogen transmission [50,51]. By comparison, young calves demonstrate a strong innate immunity to *B. bovis* that has been characterized as a Th1 response with macrophage induction of IL-12 and early IFN-γ response by NK cells [52,53]. Increasing evidence points to the central importance of macrophages, DCs and NK cells in the spleen-dependent immune response of calves to *B. bovis* [54,55]. IL-12 produced by *B. bovis* exposed monocytes was sufficient to drive IFN-γ production by NK cells [56]. Evidence of crosstalk between human DCs and NK cells has revealed that DCs can improve the effector function of NK cells and in response NK cells can promote maturation and immunostimulatory properties of DCs [57]. This interaction may be a likely target for immunosuppression by tick salivary factors. *R. sanguineus* has been shown to suppress IL-12 production in murine bone-marrow derived DCs after LPS stimulation [23]. To the best of the authors' knowledge, this report is the first to document that *R. microplus* salivary gland proteins reduce IL-12 message in bovine macrophages activated by LPS. Reduced IL-12 is consistent with lower levels of IFN-γ and TNFα, which correlate with the suppression of a Th1 response. These studies are limited to largely looking at changes in mRNA message profiles, and have not been correlated to protein levels. Additional studies are warranted to identify and characterize the immunomodulatory factors in the salivary glands of *R. microplus* and investigate the molecular mechanisms affecting bovine APCs. Examining the downstream effects on T cell memory and effector function, NK activation and crosstalk in cattle will contribute to our understanding of how components in tick saliva suppress Th1 responses. Knowledge of the immunobiological intricacies underlying the tick-host blood feeding interface offers the opportunity to innovate sustainable technologies to mitigate the impact of *R. microplus* on livestock production systems globally.

Conclusions

Molecules in the salivary glands of adult *R. microplus* showed bimodal concentration-, and time-dependent effects on differential up-regulation of CD86 in bovine macrophages activated by the TLR4-ligand, LPS. Proinflammatory cytokines and the Th1 promoting cytokine, IL-12, were down regulated in a dose-dependent manner. The co-stimulatory molecule CD80 and the activation marker CD69 were also suppressed by salivary gland extracts. Our results indicate that salivary gland factors may prevent activation of the innate immune system in parasitized cattle. Further studies to identify and characterize the immunomodulatory salivary factors secreted by *R. microplus* at the feeding site are warranted to assess their molecular mechanism, and to test

the hypothesis that they are involved in the transmission of *Babesia* spp.

Acknowledgements

Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. Our appreciation goes to Jason Tidwell for technical assistance, Keith Shelley and Larry Camarillo for animal care and sample acquisition. This study was financially supported by USDA-ARS appropriated project 6205-32000-031-00D. The USDA is an equal opportunity provider and employer.

Authors' contributions

DKB conceived, designed and performed the experiments. Manuscript was primarily written by DKB with review of the manuscript provided from AAPL. AAPL conceived, administered, and supervised funding for the experiments. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 28 September 2011 Accepted: 14 February 2012

Published: 14 February 2012

References

- Jongejan F, Uilenberg G: The global importance of ticks. *Parasitology* 2004, **129**(Suppl):S3-14.
- Bock R, Jackson L, de Vos A, Jorgensen W: Babesiosis of cattle. *Parasitology* 2004, **129**(Suppl):S247-269.
- Miller RJ, Davey RB, George JE: First report of permethrin-resistant *Boophilus microplus* (Acari: Ixodidae) collected within the United States. *J Med Entomol* 2007, **44**:308-315.
- Fernandez-Salas A, Rodriguez-Vivas RI, Alonso-Diaz MA: First report of a *Rhipicephalus microplus* tick population multi-resistant to acaricides and ivermectin in the Mexican tropics. *Vet Parasitol* 2011, **172**:109-113.
- Wikel SK: Tick modulation of host immunity: an important factor in pathogen transmission. *Int J Parasitol* 1999, **29**:851-859.
- Fontaine A, Diouf I, Bakkali N, Misse D, Pages F, Fusai T, Rogier C, Almeras L: Implication of haematophagous arthropod salivary proteins in host-vector interactions. *Parasit Vectors* 2011, **4**:187.
- Maruyama SR, Anatriello E, Anderson JM, Ribeiro JM, Brandao LG, Valenzuela JG, Ferreira BR, Garcia GR, Szabo MP, Patel S, et al: The expression of genes coding for distinct types of glycine-rich proteins varies according to the biology of three metastrongyloid ticks, *Rhipicephalus* (*Boophilus*) *microplus*, *Rhipicephalus sanguineus* and *Amblyomma cajennense*. *BMC Genomics* 2010, **11**:363.
- Murrell A, Barker SC: Synonymy of *Boophilus Curtice*, 1891 with *Rhipicephalus Koch*, 1844 (Acari: Ixodidae). *Syst Parasitol* 2003, **56**:169-172.
- Jonsson NN, Bock RE, Jorgensen WK: Productivity and health effects of anaplasmosis and babesiosis on *Bos indicus* cattle and their crosses, and the effects of differing intensity of tick control in Australia. *Vet Parasitol* 2008, **155**:1-9.
- Steen NA, Barker SC, Alewood PF: Proteins in the saliva of the Ixodida (ticks): pharmacological features and biological significance. *Toxicon* 2006, **47**:1-20.
- Brossard M, Wikel SK: Tick immunobiology. *Parasitology* 2004, **129**(Suppl): S161-176.
- Allen JR, Khalil HM, Wikel SK: Langerhans cells trap tick salivary gland antigens in tick-resistant guinea pigs. *J Immunol* 1979, **122**:563-565.
- Nithiuthai S, Allen JR: Langerhans cells present tick antigens to lymph node cells from tick-sensitized guinea-pigs. *Immunology* 1985, **55**:157-163.
- Nithiuthai S, Allen JR: Significant changes in epidermal Langerhans cells of guinea-pigs infested with ticks (*Dermacentor andersoni*). *Immunology* 1984, **51**:133-141.
- Merad M, Ginhoux F: Dendritic cell genealogy: a new stem or just another branch? *Nat Immunol* 2007, **8**:1199-1201.
- Mejri N, Brossard M: Splenic dendritic cells pulsed with *Ixodes ricinus* tick saliva prime naive CD4+T to induce Th2 cell differentiation in vitro and in vivo. *Int Immunol* 2007, **19**:535-543.
- Mejri N, Francini N, Rutti B, Brossard M: Th2 polarization of the immune response of BALB/c mice to *Ixodes ricinus* instars, importance of several antigens in activation of specific Th2 subpopulations. *Parasite Immunol* 2001, **23**:61-69.
- Muller-Doblies UU, Maxwell SS, Boppana VD, Mihalyo MA, McSorley SJ, Vella AT, Adler AJ, Wikel SK: Feeding by the tick, *Ixodes scapularis*, causes CD4(+) T cells responding to cognate antigen to develop the capacity to express IL-4. *Parasite Immunol* 2007, **29**:485-499.
- Schoeler GB, Manweiler SA, Wikel SK: Cytokine responses of C3H/HeN mice infested with *Ixodes scapularis* or *Ixodes pacificus* nymphs. *Parasite Immunol* 2000, **22**:31-40.
- Ferreira BR, Silva JS: Successive tick infestations selectively promote a T-helper 2 cytokine profile in mice. *Immunology* 1999, **96**:434-439.
- Macaluso KR, Wikel SK: *Dermacentor andersoni*: effects of repeated infestations on lymphocyte proliferation, cytokine production, and adhesion-molecule expression by BALB/c mice. *Ann Trop Med Parasitol* 2001, **95**:413-427.
- Boppana VD, Thangamani S, Alarcon-Chaidez FJ, Adler AJ, Wikel SK: Blood feeding by the Rocky Mountain spotted fever vector, *Dermacentor andersoni*, induces interleukin-4 expression by cognate antigen responding CD4+ T cells. *Parasit Vectors* 2009, **2**:47.
- Cavassani KA, Aliberti JC, Dias AR, Silva JS, Ferreira BR: Tick saliva inhibits differentiation, maturation and function of murine bone-marrow-derived dendritic cells. *Immunology* 2005, **114**:235-245.
- Vesely DL, Fish D, Shlomchik MJ, Kaplan DH, Bockenstedt LK: Langerhans cell deficiency impairs *Ixodes scapularis* suppression of Th1 responses in mice. *Infect Immun* 2009, **77**:1881-1887.
- Sa-Nunes A, Bafica A, Lucas DA, Conrads TP, Veenstra TD, Andersen JF, Mather TN, Ribeiro JM, Francischetti IM: Prostaglandin E2 is a major inhibitor of dendritic cell maturation and function in *Ixodes scapularis* saliva. *J Immunol* 2007, **179**:1497-1505.
- Hajnicka V, Vancova I, Kocakova P, Slovak M, Gasperik J, Slavikova M, Hails RS, Labuda M, Nuttall PA: Manipulation of host cytokine network by ticks: a potential gateway for pathogen transmission. *Parasitology* 2005, **130**:333-342.
- Sharpe AH, Freeman GJ: The B7-CD28 superfamily. *Nat Rev Immunol* 2002, **2**:116-126.
- Chen XQ, Yang J, Hu SP, Nie HX, Mao GY, Chen HB: Increased expression of CD86 and reduced production of IL-12 and IL-10 by monocyte-derived dendritic cells from allergic asthmatics and their effects on Th1 and Th2-type cytokine balance. *Respiration* 2006, **73**:34-40.
- Ranger AM, Das MP, Kuchroo VK, Glimcher LH: B7-2 (CD86) is essential for the development of IL-4-producing T cells. *Int Immunol* 1996, **8**:1549-1560.
- Yoshida T, Hachimura S, Ishimori M, Ise W, Totsuka M, Ametani A, Kaminogawa S: Interleukin 12 and CD86 Regulate Th1 and Th2 Development Induced by a Range of Antigen Doses Presented by Peyer's Patch and Spleen Cells. *Cytotechnology* 2003, **43**:81-88.
- Brown JA, Titus RG, Nabavi N, Glimcher LH: Blockade of CD86 ameliorates *Leishmania major* infection by down-regulating the Th2 response. *J Infect Dis* 1996, **174**:1303-1308.
- Wang Z, Takamoto M, Sugane K: Costimulatory signal through CD86 is important in Th2 response in *Trichinella spiralis* infected mice. *Parasite Immunol* 2000, **22**:121-130.
- Deszo EL, Brake DK, Kelley KW, Freund GG: IL-4-dependent CD86 expression requires JAK/STAT6 activation and is negatively regulated by PKCdelta. *Cell Signal* 2004, **16**:271-280.
- Brake DK, Wikel SK, Tidwell JP, Perez de Leon AA: *Rhipicephalus microplus* salivary gland molecules induce differential CD86 expression in murine macrophages. *Parasit Vectors* 2010, **3**:103.
- Guerrero FD, Bendele KG, Davey RB, George JE: Detection of *Babesia bigemina* infection in strains of *Rhipicephalus* (*Boophilus*) *microplus* collected from outbreaks in south Texas. *Vet Parasitol* 2007, **145**:156-163.
- Raaijmakers MH, van Ernt L, de Witte T, Mensink E, Raymakers RA: Quantitative assessment of gene expression in highly purified hematopoietic cells using real-time reverse transcriptase polymerase chain reaction. *Exp Hematol* 2002, **30**:481-487.
- Inokuma H, Kerlin RL, Kemp DH, Willadsen P: Effects of cattle tick (*Boophilus microplus*) infestation on the bovine immune system. *Vet Parasitol* 1993, **47**:107-118.

38. Piper EK, Jackson LA, Bagnall NH, Kongsuwan KK, Lew AE, Jonsson NN: **Gene expression in the skin of *Bos taurus* and *Bos indicus* cattle infested with the cattle tick, *Rhipicephalus (Boophilus) microplus*.** *Vet Immunol Immunopathol* 2008, **126**:110-119.
39. Piper EK, Jonsson NN, Gondro C, Lew-Tabor AE, Moolhuijzen P, Vance ME, Jackson LA: **Immunological profiles of *Bos taurus* and *Bos indicus* cattle infested with the cattle tick, *Rhipicephalus (Boophilus) microplus*.** *Clin Vaccine Immunol* 2009, **16**:1074-1086.
40. Carvalho WA, Franzin AM, Abatepaulo AR, de Oliveira CJ, More DD, da Silva JS, Ferreira BR, de Miranda Santos IK: **Modulation of cutaneous inflammation induced by ticks in contrasting phenotypes of infestation in bovines.** *Vet Parasitol* 2010, **167**:260-273.
41. Carvalho WA, Maruyama SR, Franzin AM, Abatepaulo AR, Anderson JM, Ferreira BR, Ribeiro JM, More DD, Augusto Mendes Maia A, Valenzuela JG, et al: ***Rhipicephalus (Boophilus) microplus*: clotting time in tick-infested skin varies according to local inflammation and gene expression patterns in tick salivary glands.** *Exp Parasitol* 2010, **124**:428-435.
42. Kashino SS, Resende J, Sacco AM, Rocha C, Proenca L, Carvalho WA, Firmino AA, Queiroz R, Benavides M, Gershwil LJ, De Miranda Santos IK: ***Boophilus microplus*: the pattern of bovine immunoglobulin isotype responses to high and low tick infestations.** *Exp Parasitol* 2005, **110**:12-21.
43. Jeannin P, Delneste Y, Lecoanet-Henchoz S, Gauchat JF, Ellis J, Bonnefoy JY: **CD86 (B7-2) on human B cells. A functional role in proliferation and selective differentiation into IgE- and IgG4-producing cells.** *J Biol Chem* 1997, **272**:15613-15619.
44. Anatriello E, Ribeiro JM, de Miranda-Santos IK, Brandao LG, Anderson JM, Valenzuela JG, Maruyama SR, Silva JS, Ferreira BR: **An insight into the sialotranscriptome of the brown dog tick, *Rhipicephalus sanguineus*.** *BMC Genomics* 2010, **11**:450.
45. Prates DB, Santos LD, Miranda JC, Souza AP, Palma MS, Barral-Netto M, Barral A: **Changes in amounts of total salivary gland proteins of *Lutzomyia longipalpis* (Diptera: Psychodidae) according to age and diet.** *J Med Entomol* 2008, **45**:409-413.
46. Leboulle G, Rochez C, Louahed J, Ruti B, Brossard M, Bollen A, Godfroid E: **Isolation of *Ixodes ricinus* salivary gland mRNA encoding factors induced during blood feeding.** *Am J Trop Med Hyg* 2002, **66**:225-233.
47. Testi R, D'Ambrosio D, De Maria R, Santoni A: **The CD69 receptor: a multipurpose cell-surface trigger for hematopoietic cells.** *Immunol Today* 1994, **15**:479-483.
48. Sancho D, Gomez M, Sanchez-Madrid F: **CD69 is an immunoregulatory molecule induced following activation.** *Trends Immunol* 2005, **26**:136-140.
49. Martin P, Gomez M, Lamana A, Matesanz Marin A, Cortes JR, Ramirez-Huesca M, Barreiro O, Lopez-Romero P, Gutierrez-Vazquez C, de la Fuente H, et al: **The leukocyte activation antigen CD69 limits allergic asthma and skin contact hypersensitivity.** *J Allergy Clin Immunol* 2010, **126**:355-365, 365 e351-353.
50. Zeidner NS, Dolan MC, Massung R, Piesman J, Fish D: **Coinfection with *Borrelia burgdorferi* and the agent of human granulocytic ehrlichiosis suppresses IL-2 and IFN gamma production and promotes an IL-4 response in C3H/HeJ mice.** *Parasite Immunol* 2000, **22**:581-588.
51. Zeidner NS, Schneider BS, Rutherford JS, Dolan MC: **Suppression of Th2 cytokines reduces tick-transmitted *Borrelia burgdorferi* load in mice.** *J Parasitol* 2008, **94**:767-769.
52. Goff WL, Johnson WC, Parish SM, Barrington GM, Tuo W, Valdez RA: **The age-related immunity in cattle to *Babesia bovis* infection involves the rapid induction of interleukin-12, interferon-gamma and inducible nitric oxide synthase mRNA expression in the spleen.** *Parasite Immunol* 2001, **23**:463-471.
53. Goff WL, Johnson WC, Horn RH, Barrington GM, Knowles DP: **The innate immune response in calves to *Boophilus microplus* tick transmitted *Babesia bovis* involves type-1 cytokine induction and NK-like cells in the spleen.** *Parasite Immunol* 2003, **25**:185-188.
54. Schneider DA, Yan H, Bastos RG, Johnson WC, Gavin PR, Allen AJ, Barrington GM, Herrmann-Hoesing LM, Knowles DP, Goff WL: **Dynamics of bovine spleen cell populations during the acute response to *Babesia bovis* infection: an immunohistological study.** *Parasite Immunol* 2010, **33**:34-44.
55. Brown WC, Norimine J, Knowles DP, Goff WL: **Immune control of *Babesia bovis* infection.** *Vet Parasitol* 2006, **138**:75-87.
56. Goff WL, Storset AK, Johnson WC, Brown WC: **Bovine splenic NK cells synthesize IFN-gamma in response to IL-12-containing supernatants**

- from *Babesia bovis*-exposed monocyte cultures. *Parasite Immunol* 2006, **28**:221-228.
57. Wehner R, Dietze K, Bachmann M, Schmitz M: **The bidirectional crosstalk between human dendritic cells and natural killer cells.** *J Innate Immun* 2011, **3**:258-263.

doi:10.1186/1756-3305-5-38

Cite this article as: Brake and Pérez de León: Immunoregulation of bovine macrophages by factors in the salivary glands of *Rhipicephalus microplus*. *Parasites & Vectors* 2012 **5**:38.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

