

RESEARCH

Open Access

# Systematics of the Oswaldoi Complex (*Anopheles*, *Nyssorhynchus*) in South America

Freddy Ruiz-Lopez<sup>1,2\*</sup>, Richard C Wilkerson<sup>1,3</sup>, David J Ponsonby<sup>4</sup>, Manuela Herrera<sup>5</sup>, Maria Anice Mureb Sallum<sup>6</sup>, Ivan Dario Velez<sup>2</sup>, Martha L Quiñones<sup>5</sup>, Carmen Flores-Mendoza<sup>7</sup>, Dave D Chadee<sup>8</sup>, Joubert Alarcon<sup>9</sup>, Joubert Alarcon-Ormasa<sup>9</sup> and Yvonne-Marie Linton<sup>3,10</sup>

## Abstract

**Background:** Effective malaria control relies on accurate identification of those *Anopheles* mosquitoes responsible for the transmission of *Plasmodium* parasites. *Anopheles oswaldoi* s.l. has been incriminated as a malaria vector in Colombia and some localities in Brazil, but not ubiquitously throughout its Neotropical range. This evidence together with variable morphological characters and genetic differences supports that *An. oswaldoi* s.l. comprises a species complex. The recent fully integrated redescription of *An. oswaldoi* s.s. provides a solid taxonomic foundation from which to molecularly determine other members of the complex.

**Methods:** DNA sequences of the Second Internal Transcribed Spacer (ITS2 - rDNA) (n = 192) and the barcoding region of the *Cytochrome Oxidase I* gene (*COI* - mtDNA) (n = 110) were generated from 255 specimens of *An. oswaldoi* s.l. from 33 localities: Brazil (8 localities, including the lectotype series of *An. oswaldoi*), Ecuador (4), Colombia (17), Trinidad and Tobago (1), and Peru (3). *COI* sequences were analyzed employing the Kimura-two-parameter model (K2P), Bayesian analysis (MrBayes), Mixed Yule-Coalescent model (MYC, for delimitation of clusters) and TCS genealogies.

**Results:** Separate and combined analysis of the *COI* and ITS2 data sets unequivocally supported four separate species: two previously determined (*An. oswaldoi* s.s. and *An. oswaldoi* B) and two newly designated species in the Oswaldoi Complex (*An. oswaldoi* A and *An. sp. nr. konderi*). The *COI* intra- and inter-specific genetic distances for the four taxa were non-overlapping, averaging 0.012 (0.007 to 0.020) and 0.052 (0.038 to 0.064), respectively. The concurring four clusters delineated by MrBayes and MYC, and four independent TCS networks, strongly confirmed their separate species status. In addition, *An. konderi* of Sallum should be regarded as unique with respect to the above. Despite initially being included as an outgroup taxon, this species falls well within the examined taxa, suggesting a combined analysis of these taxa would be most appropriate.

**Conclusions:** Through novel data and retrospective comparison of available *COI* and ITS2 DNA sequences, evidence is shown to support the separate species status of *An. oswaldoi* s.s., *An. oswaldoi* A and *An. oswaldoi* B, and at least two species in the closely related *An. konderi* complex (*An. sp. nr. konderi*, *An. konderi* of Sallum). Although *An. oswaldoi* s.s. has never been implicated in malaria transmission, *An. oswaldoi* B is a confirmed vector and the new species *An. oswaldoi* A and *An. sp. nr. konderi* are circumstantially implicated, most likely acting as secondary vectors.

**Keywords:** *Anopheles oswaldoi* species complex, *An. oswaldoi* s.s., *An. oswaldoi* A, *An. oswaldoi* B, *An. sp. nr. konderi*, *COI* barcoding, ITS2

\* Correspondence: fredyruiz9@gmail.com

<sup>1</sup>Department of Entomology, National Museum of Natural History, Smithsonian Institution, Museum Support Center, Suitland, MD 20746, USA

<sup>2</sup>Programa de Estudio y Control de Enfermedades Tropicales (PECET),

Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia

Full list of author information is available at the end of the article

## Background

Species complexes are relatively common in the family Culicidae [1], and several Neotropical *Anopheles*, including some vector species, are known to comprise isomorphic species. Within the Oswaldoi Group alone (*Anopheles*, subgenus *Nyssorhynchus*), seven of the 15 formally recognised species comprise complexes [2-11]. *Anopheles oswaldoi* (Peryassú) is one such taxon. It is thought to comprise a species complex in Brazil [4,9,12-16] and Colombia [8,13,17] and has been implicated in malaria transmission in some parts of its range [11-13], yet its taxonomic and vectorial status elsewhere in South America remains unclear. A comprehensive revision of the taxonomy and current distribution of *An. oswaldoi* is given in Motoki *et al.* [12].

Several studies have provided evidence for genetic variation in *An. oswaldoi*. Firstly, Marrelli *et al.* [14] analyzed ITS2 sequences from seven populations of *An. oswaldoi* s.l. and determined four geographic groups, as follows: Group I from Acre, Amazonas and Rondônia (Brazil), Group II from Ocamo (Venezuela) and Amapá (Brazil), Group III from Espírito Santo (type locality), Brazil and Group IV from Yurimaguas, Peru. Subsequently, Ruiz *et al.* [17] revised these groupings and determined that Group II corresponded to *An. oswaldoi* B from Putumayo, Colombia [13], and that Group IV from Yurimaguas, Peru, was actually *An. benarrochi* B [17], a newly recognized species of the subgenus *Nyssorhynchus* [17,18]. The identification of Group III (GenBank: AF055072) from the type locality of Espírito Santo in Brazil was later corrected to that of *An. evansae* (Bréthes) [4]. The misidentification resulted from the incorrect use of polymorphic characters in the wing (humeral pale spot) and second hindtarsal segment (basal dark band); characters that overlap between *An. oswaldoi* and *An. evansae* in currently available taxonomic keys [11]. The true identity of Group I remains unclear.

Scarpassa and Conn [16] sequenced a long fragment of *COI* from 45 *An. oswaldoi* s.l. from four populations from Brazil (Acre, Amazonas, Rondônia and Pará). Parsimony analysis revealed four distinct groups: Group I, Acre (Sena Madureira) and Rondônia (São Miguel); Group II, Rondônia (São Miguel); Group III, Pará (Moju), and Group IV from Acre (Sena Madureira) and Coari (Amazonas). Although the authors tentatively suggested that Group I may be *An. oswaldoi* s.s. and Group IV may be *An. konderi* Galvão and Damasceno, based primarily on geographic origin, they lacked certainty in assigning taxonomic names to these phylogenetic lineages.

The taxonomic confusion between *An. oswaldoi* and *An. konderi* is not new. *Anopheles konderi* was originally described from specimens collected in the Solimões River at Coari, Amazonas, Brazil in 1942 [19]. Soon after, Lane [20] reduced it to a junior synonym of *An. oswaldoi*, where

it remained until its re-elevation to separate species status in 2004 [21]. Isoenzyme analysis of 20 loci in three populations of purported *An. oswaldoi* and *An. konderi* from the Brazilian Amazon (Coari, Amazonas (*An. konderi*); São Miguel, Rondônia (*An. oswaldoi* and *An. konderi*) and Sena Madureira, Acre (*An. oswaldoi* and *An. konderi*)) revealed no significant differences between populations, and led the author to question whether *An. oswaldoi* and *An. konderi* were indeed truly separate species, or whether ongoing introgression between the two species would explain this low level of variation [22]. Recently, Motoki *et al.* [23] using *COI*, *white* and ITS2 DNA sequences confirmed that morphologically identified specimens of *An. konderi* comprised at least three species in the Amazonian region. Unfortunately, none of these recent studies (including the redescription [21]) have examined specimens from the type locality, thus the true identity of *An. konderi* s.s. remains unclear.

The same was true for *An. oswaldoi* s.s. until a recent redescription of the species was undertaken based on progeny broods collected from the type locality of Espírito Santo, Brazil [12]. This study included morphological data for immature stages (fourth-instar larvae and pupae) and adults of both sexes, as well as corresponding ITS2 DNA sequence. As a holotype was not designated for *An. oswaldoi* s.s. when it was first described by Peryassú in 1922, a lectotype was chosen from the type series [12]. This study provided a solid taxonomic platform from which to attempt to further determine the component members of the Oswaldoi Complex. Comparison of DNA sequences of *An. oswaldoi* s.s. [12] confirms that neither Marrelli *et al.* [14] (ITS2), nor Scarpassa and Conn [16] (*COI*) included “true” *An. oswaldoi* in their studies. To date, *An. oswaldoi* s.s. has only been confirmed in the Brazilian States of Espírito Santo, Rio de Janeiro and São Paulo in Brazil [12,24]. Pinault and Hunter [25] recently reported three *COI* sequences of *An. oswaldoi* from Ecuador (GenBank: JN412831-33), however these sequences were misidentified, sharing high similarity with those of *An. rangeli* (GenBank: HM022390-94). Based on morphology and ITS2 sequences, Sallum *et al.* [9] detected at least two cryptic species in *An. oswaldoi* s.l. collected in the state of Acre (Brazil), both of which differ from *An. oswaldoi* s.s.

Given the evidence above, there is no doubt that *An. oswaldoi* comprises a species complex in Latin America. The objectives of this study were to ascertain the taxonomic status and relative distribution of the component members of the *An. oswaldoi* species complex in Brazil, Colombia, Ecuador, Peru, Trinidad and Tobago and Venezuela using ITS2 and *COI* barcodes, and to correlate this information with previously documented vector incrimination studies of *An. oswaldoi* s.l. across its range.

## Methods

### Specimens

A total of 255 specimens of *Anopheles oswaldoi* s.l. from 33 localities in five countries in the Neotropics (Brazil, Colombia, Ecuador, Peru and Trinidad and Tobago) were used in this study (see Table 1 for georeferenced locality data). Co-ordinates were converted to decimal degrees [26] and the distribution data is available in Mosquito Map ([www.mosquitomap.org](http://www.mosquitomap.org)). Samples of *Anopheles konderi* s.l. were collected near Macapá, Amapá, Brazil and identified

on the basis of the male aedeagus by MAMS. To avoid confusion, these are referred to herein as *An. konderi* of Sallum. All specimens used in this study were collected or provided by the authors of this study.

### Molecular analysis

DNA was extracted using the DNeasy® Blood and Tissue Kit (QIAGEN®, USA) on the automated BioSprint 96® robotic platform. The ITS2 region was amplified for 192 samples from 21 localities using the published primers

**Table 1 Origin and georeferences of *An. oswaldoi* s.l. specimens used in this study, showing relative numbers of COI (n = 110) and ITS2 (n = 192) sequences obtained from 255 specimens have been used**

Country	State and exact locality	n =	COI	ITS2 (H)	Latitude	Longitude
Brazil	Amazonas, Tefé	1	1	1 (III)	-03.3207	-64.7236
	Espirito Santo, Jaguaré, Fazenda Marianelli	10	3	10 (I)	-19.0348	-39.9485
	Mato Grosso, Peixoto de Azevedo	42	42	14 (II, IV-IX)	-10.2257	-54.9862
	Rio de Janeiro, Morro de Panela	2	2	-	-22.9678	-43.3-415
	Rondônia, Ariquemes	3	3	-	-09.9136	-63.0440
	Rondônia, Costa Marques	3	3	-	-12.4156	-64.2215
	São Paulo, Pariquera-Açu	2	2	2 (I)	-24.9875	-47.9561
	São Paulo, Pariquera-Açu	1	1	-	-24.7096	-47.8839
Colombia	Amazonas, Kilometro 12	1	1	1 (III)	-04.1159	-69.9522
	Antioquia, Nechí, Mala Noche	28	12	28 (X)	08.1101	-74.7671
	Caquetá, Peñas Coloradas	2	2	1 (XII)	00.8699	-73.8419
	Meta, Granada, Morichal	2	2	-	03.5372	-73.7009
	Norte de Santander, Tibú	1	1	1 (X)	08.6403	-72.7371
	Putumayo, Agua Negra	14	-	14 (XII)	00.7494	-75.3834
	Putumayo, Cecilia Cocha	5	-	5 (XII)	00.1158	-74.9781
	Putumayo, La Apaya	6	-	6 (XII)	00.7494	-75.3833
	Putumayo, Pto. Asís, Cecilia Cocha	24	-	24 (XII)	00.1158	-74.9781
	Putumayo, Pto. Asís, La Manuela	4	4	-	00.5133	-76.4992
	Putumayo, Pto. Leguizamo, Bella Vista	8	-	8 (XII)	00.7494	-75.3833
	Putumayo, Pto. Leguizamo, El Salado	9	-	9 (XII)	00.2108	-74.8036
	Putumayo, Pto. Leguizamo, La Quebradita	4	4	2 (XII)	00.5133	-76.4992
	Putumayo, Pto. Leguizamo, Lagarto Cocha	39	-	39 (XII)	00.2108	-74.8036
	Putumayo, Pto. Leguizamo, Puntales	13	-	13 (XII)	00.4272	-74.3986
	Putumayo, Pto. Leguizamo, Tukare	1	1	1 (XII)	00.5133	-76.4992
Putumayo, Pto. Nariño	2	-	2 (XII)	00.7494	-75.3833	
Ecuador	Orellana, Coca, Cañon de los Monos	2	2	-	-00.3434	-77.0070
	Orellana, Coca, Guamayacu	4	4	-	00.1300	-77.2313
	Orellana, Coca, Juan Montalvo	7	7	-	-00.4725	-76.9914
	Orellana, Tiputini	2	2	-	-00.6381	-76.1450
Peru	Loreto, Iquitos	1	1	-	-03.7561	-73.2706
	Loreto, Rio Putumayo	1	1	-	-04.2325	-74.2179
	Madre de Dios, Davila	1	1	1 (XIII)	-11.7669	-70.8119
Trinidad and Tobago	Valencia, St. Andrew/St. David	10	8	10 (XI)	10.6447	-61.0936

Sequences generated per country include: Brazil (27 ITS2, 57 COI), Colombia (154 ITS2, 27 COI), Ecuador (15 COI), Peru (1 ITS2, 3 COI) and Trinidad and Tobago (10 ITS2, 8 COI). H: ITS2 haplotype number.

of Collins and Paskewitz [27] and following the protocol in Linton *et al.* [28]. DNA barcodes [29] were amplified from 110 individuals from 24 localities using the universal barcoding primers developed by Folmer *et al.* [30] and the protocol of the Mosquito Barcoding Initiative, expressly listed in Ruiz *et al.* [31].

Sequencing reactions were carried out in both directions using the Big Dye Terminator Kit® on an ABI3770 automated sequencer (PE Applied BioSystems®, Warrington, England). Sequence chromatograms were edited using Sequencher™ v. 4.8 (Genes Codes Corporation®, Ann Arbor, MI). Sequences were aligned automatically in Mafft Pro 5.5.7 ([www.geneious.com](http://www.geneious.com)) or using MacClade v. 4.06 [32]. Basic Local Alignment Search Tool (BLAST) searches (<http://blast.ncbi.nlm.nih.gov>) were carried out to correlate our sequences with those publicly available in GenBank.

To assess population-level genealogies, *COI* sequences were analyzed using TCS v. 1.21 [33]. A connection limit of 95 % was adopted to investigate whether *An. oswaldoi* formed a single “meta-population” (reflected by a single network) or is comprised of separate species in South America (reflected by the formation of two or more independent networks).

A *COI* data matrix was generated in MEGA v. 5 [34]. *COI* sequences were grouped according to the results of the TCS analysis, and intra- and inter-group genetic distances compared using Kimura’s 2-Parameter distance (K2P) algorithm [35]. Phylogenetic analysis was carried out on the separate and combined ITS2 and *COI* data sets. MrModeltest v. 2.3 [36] was used to choose the best evolutionary model for these regions separately using the Akaike Information Criterion (AIC) search. A partitioned Bayesian analysis (by DNA region) was subsequently performed using MrBayes v. 3.1.2 [37], available online (<http://cbsuapps.tc.cornell.edu/mrbayes.aspx>).

The analysis in MrBayes ran for 10 million generations with two parallel searches using three heated and one cold Markov chain. The first 5 million generations were discarded as burn-in. Support for this Bayesian tree was conducted by generating a maximum parsimony (MP) tree in PAUP v. 4.0b10 [38] for both ITS2 and *COI*, and bootstrapping [39], using a heuristic search, simple step-wise addition, TBR branch swapping and 1000 bootstrap replicates.

Unique *COI* haplotypes were further analysed using MYC [40,41] for delimitation of *COI* clusters. This method optimizes a threshold age that corresponds to the shift from coalescent to species diversification [42] branching processes and calculates the number of resulting independent entities. The likelihood of the null model that all samples belong to a single species is compared to that of the alternative hypothesis where separate coalescent

groups are nested within the species tree. Confidence limits correspond to threshold values  $\pm 2 \log L$  units around the ML estimate. This analysis was conducted on Bayesian consensus trees and each tree was converted to ultrametric using penalized likelihood as implemented in r8s v.1.7 [43], with the optimal smoothing parameter selected by cross-validation of values between 0.01 and 1000. FigTree v. 1.2.1 [44] was used to edit all trees generated.

## Results


Specimens of *An. oswaldoi s.l.* ( $n = 255$ ) were obtained from 33 localities in five South American countries as follow (Table 1): Brazil ( $n = 64$ ), Colombia ( $n = 163$ ), Ecuador ( $n = 15$ ), Peru ( $n = 3$ ) and Trinidad and Tobago ( $n = 10$ ). From these mosquitoes, 192 ITS2 sequences were generated and a subset of 110 of these, including four specimens from the redescription [12], were also barcoded. The full data sets by country follow: Brazil (27 ITS2; 57 *COI*, GenBank: KF809034-078, KF809121-132), Colombia (154 ITS2; 27 *COI*, GenBank: KF809079, KF809085-091, KF809093-096, KF809100, KF809102, KF809105, KF809109-119, KF809133), Ecuador (15 *COI*, GenBank: KF809081-084, KF809099, KF809103, KF809108, KF809120, KF809135-139, KF809142-143), Peru (1 ITS2; 3 *COI*, GenBank: KF809134, KF809140-141) and Trinidad and Tobago (10 ITS2; 8 *COI*, GenBank: KF809080, KF809092, KF809097-098, KF809101, KF809104, KF809106-107, KF809128). *COI* sequences of *An. konderi sensu Sallum* ( $n=4$ ) are available in GenBank under accessions KF809030-033.

### ITS2 sequences of *An. oswaldoi s.l.*

ITS2 sequences were generated for 192 specimens of *An. oswaldoi s.l.* (Table 1). The overall alignment (544 bp, after primer trim) revealed 13 unique ITS2 haplotypes (GenBank: KC970065-77), labelled HI-HXIII (Table 2). The ITS2 sequences ranged in length from 530 bp in Haplotype I (H1) (Espírito Santo and São Paulo) to 540 bp in haplotypes HVI-IX from Mato Grosso in Brazil (Table 2). Thirty-two variable bases were noted (5.9 %), with the variation concentrated toward the second half (3’ end) of the ITS2 fragment between aligned nucleotides 207–494. Indels (insertions/deletions) were observed at bases 337–338, 344–345, 365–369, 408–409, 416–419, and 467–468 (Table 2).

Whereas in general all geographical areas were represented by unique ITS2 haplotypes, the 14 specimens of *An. oswaldoi* from Mato Grosso (Brazil) showed seven unique haplotypes (HII, HIV - HIX), varying in length between 538 bp and 540 bp (Table 2). Specimens from Putumayo, Colombia ( $n = 123$ ) and Caquetá, Colombia ( $n = 1$ ) shared the same ITS2 haplotype (HXII) and one specimen from Amazonas, Colombia shared the same





sp. nr. *konderi*) (Figure 3). The *COI* sequence of *An. konderi* of Sallum was found to cluster closely with *An. sp. nr. konderi*, but these two species are consistently different in four of their ITS2 bases as follows: one transition in position 361 (G/A), two indels (364 (A/-) and 365 (T/-)), and one transversion (449 T/C) (data not shown).

Results of the TCS, MYC model and Bayesian analyses using the ITS2 and *COI* data sets provides further strong

support for the following species: *An. oswaldoi* s.s. from Brazil (Espírito Santo = type locality), Rondônia, Rio de Janeiro and São Paulo); *An. oswaldoi* A from Brazil (Mato Grosso, Rondônia, Amazonas) and Colombia (Amazonas); *An. oswaldoi* B from Colombia (Antioquia, Caquetá, Meta, Norte de Santander, Putumayo), Ecuador (Orellana) and Trinidad and Tobago (Valencia); and *An. sp. nr. konderi* from Colombia (Caquetá), Ecuador (Orellana) and Peru (Madre de Dios, Loreto) (Figures 1, 2, 3 and 4).

**Table 3** Mean inter- and intra-specific pairwise distances (K2P model)


Species	n =	osw s.s.	osw A	osw B	sp. nr. kon
<i>An. oswaldoi</i> s.s.	12	<b>0.007</b>			
<i>An. oswaldoi</i> A	46	0.057	<b>0.012</b>		
<i>An. oswaldoi</i> B	41	0.038	0.064	<b>0.020</b>	
<i>An. sp. nr. konderi</i>	11	0.045	0.053	0.056	<b>0.010</b>

Analyses of *COI* sequences separated 110 specimens of *Anopheles oswaldoi* s.l. into four groups using the TCS model as calculated by Kimura's 2-Parameter distance model (K2P) [35].

## Discussion

### *COI* and ITS2 sequence analyses

Based on mitochondrial *COI* and nuclear ITS2 sequences, this study presents clear molecular evidence that the Oswaldoi-Konderi Complex comprises at least five species: two previously determined (*An. oswaldoi* s.s. and *An. oswaldoi* B), as well as *An. oswaldoi* A and *An. sp. nr. konderi*, and *An. konderi* of Sallum defined herein for the first time. Phylogenetic reconstruction of *COI*




and ITS2 sequences (both independently and concatenated) verified four distinct species, which concur with the MYC and TCS network models. Construction of genealogical networks using TCS are often used to infer evolutionary relationships at the population level; however, identification of independent networks that fall beyond the 95% parsimony connect limit are considered putative biological species [45]. At this threshold, four independent networks were generated corresponding to the four species listed above (Figure 1). In addition samples of *An. konderi* of Sallum, initially included as an outgroup in this study, not only represented a separate taxon but also fell firmly within the Oswaldoi Complex in phylogenetic treatments. This supports that *An. oswaldoi* and *An. konderi* comprise species complexes of very closely related taxa, which would benefit from being assessed concurrently.

Following this robust molecular delimitation, efforts were made to retrospectively identify samples documented in previously published studies, and relate these to geographical distributions and their regional malaria vector status.

#### *Anopheles oswaldoi* s.s.

Prior to this study, the distribution of “true” *An. oswaldoi* s.s. had only been determined from Espírito Santo and São Paulo, Brazil [12]. Herein *COI* data also confirms its presence in the Brazilian States of Rio de Janeiro, Acre (GenBank: DQ784832-35) and Coari (GenBank: DQ784836-38) [15] (Table 4). Ruiz *et al.* [8,31] suggested a genetic distance threshold of 0.02 for species delimitation in *Nyssorhynchus* species. Despite the large geographic distance, our *An. oswaldoi* s.s. from São Paulo and Espírito Santo, Brazil (type locality, south of Brazil) grouped with Scarpassa and Conn’s *COI* sequences [16] from Rondônia State, in the north of Brazil (overall mean K2P distance of 0.010, range 0.002 to 0.015). Given its current distribution, it seems highly likely that *An. oswaldoi* s.s. could also be present in regions between these Brazilian states, including Mato Grosso, Goiás and Minas Gerais, and perhaps even south into Bolivia or Paraguay.

There are no records to suggest that *An. oswaldoi* s.s. is a vector in the eastern Brazilian provinces of Espírito Santo, Rio de Janeiro or São Paulo. In the north-western




**Figure 3** Combined MrBayes tree of *An. oswaldoi* s.l. using ITS2 and COI sequences. Four groups are clearly defined *An. oswaldoi* s.s. (Espírito Santo, Rio de Janeiro, Rondônia and São Paulo, Brazil); *An. oswaldoi* A (Mato Grosso, Rondônia and Tefé, Brazil and Amazonas, Colombia); *An. oswaldoi* B from Colombia (Antioquia, Caquetá, Meta, Norte de Santander and Putumayo), Ecuador (Orellana) and Trinidad and Tobago (Valencia); and *An. sp. nr. konderi* from Ecuador (Orellana), Colombia (Caquetá) and Peru (Loreto and Madre de Dios). B: Bootstrap; P: Posterior probability. *Anopheles konderi* of Sallum from near Macapá, Amapá, Brazil, is clearly differentiated from the other species of the *An. oswaldoi* complex, and appears to sit within the Oswaldoi Group. \*Outgroups: *An. albitarsis* F and *An. nuneztovari* C.

Brazilian state of Rondônia, where *An. oswaldoi* s.s., *An. oswaldoi* A and *An. sp. nr. konderi* are present, de Oliveira-Ferreira *et al.* [46] and Klein *et al.* [47,48] reported low levels of *Plasmodium* infections in what can only be assumed to include a mix of *oswaldoi-konderi* lineages.

#### ***Anopheles oswaldoi* B**

In this study, three ITS2 haplotypes [HX, HXI and HXII (Table 2)] were found to be identical to those reported as *An. oswaldoi* B from Putumayo, Colombia by Ruiz *et al.* [17,31]. These haplotypes shared 99% identity to published sequences from Ocamo, Venezuela (GenBank: AF055071)





and Amapá, Brazil (GenBank: AF056318) (= Group II of Marrelli *et al.* [14]). Our analyses of both individual and combined COI and ITS2 data sets, grouped the above sequences into a single phylogenetic cluster and TCS network, confirming *An. oswaldoi* B as a separate species in the *An. oswaldoi* complex. This study increases the known distribution of *An. oswaldoi* B to include the Departments of Antioquia, Caquetá and Norte de Santander (Colombia), and the Province of Orellana (Ecuador) and Saint Andrew/Saint David (Trinidad and Tobago) (Figure 4).

*Plasmodium vivax* infection was detected in one molecularly confirmed *An. oswaldoi* B specimen (of 361 positive mosquitoes) collected in Putumayo, Colombia [13]. However *An. oswaldoi* s.l. has also been incriminated in Amapá, Brazil [49], as a secondary vector in Venezuela [50] and has been experimentally infected with *P. vivax* in Trinidad and Tobago [51], where only *An. oswaldoi* B has been identified to date. Correlation of these studies with the known distribution of *An. oswaldoi* B suggest the potential involvement of this species on malaria transmission is over a much wider region of northern and northeastern South America than originally realised.

Currently, the only valid synonym of *An. oswaldoi* is *An. aquacaelestis* Curry, originally described from Panama [52].

The original description differentiates *An. aquacaelestis* from *An. aquasalis*, and suggests that this Panamanian species is identical to the species present in Trinidad and Tobago. As *An. oswaldoi* B is the only species of the *An. oswaldoi* complex that has been detected in Trinidad and Tobago, it would be prudent to carefully study the type series of *An. aquacaelestis*, to ensure that our informally designated "*An. oswaldoi* B" is not in fact *An. aquacaelestis*, prior to assigning a new name.

#### *Anopheles oswaldoi* A (designated herein)

Comparison of our *An. oswaldoi* A ITS2 sequences (as haplotypes HIII-IX; Table 2) revealed high similarity with three in Marrelli *et al.* [4] and two in Sallum *et al.* [9]. Our Haplotype III from Tefé, Amazonas, Brazil and Amazon, Colombia, shares 99.8% identity with GenBank AF056317 from Amazonas, Brazil (= Group I of Marrelli *et al.* [14]) and varies by only a single base indel (-/T) at position 166. Our ITS2 haplotype HII from Mato Grosso, Brazil, is 99.4% identical to GenBank AF055068 from Acre, Brazil (= Group I of Marrelli *et al.* [14]) with base substitutions at positions 349 (C/A) and 427 (G/A). Our ITS2 haplotype HVI from Mato Grosso, Brazil, is 98.6% similar to GenBank AF055069 from Rondônia, Brazil (= Group I of Marrelli *et al.* [14]), and differs at only one base (position 370; G/A) from EU636802 [9]. Sallum *et al.* [9] described morphological differences in the male genitalia and ITS2 sequences of *An. oswaldoi* s.l. from Acrelândia, Acre, Brazil, compared to *An. oswaldoi* s.s. [12]. Our ITS2 haplotype HIV from Mato Grosso is identical to GenBank EU636809 from Acrelândia, Acre. These results clearly show that the unnamed species of *An. oswaldoi* s.l. described by Marrelli *et al.* [14] and Sallum *et al.* [9] corresponds to our *An. oswaldoi* A.

The COI sequences generated by Scarpassa and Conn [16] overlapped our barcode region by 404 bp. Comparison of their trimmed sequences with our data, revealed sequences from their "Group 3" specimens (GenBank DQ784849-51) from Moju, Pará, Brazil were 99% identical to those of our *An. oswaldoi* A, thus confirming that *An. oswaldoi* A is also present in Pará State (Table 4).

Drawing from our data and correlation with other published works [9,14,16], the known distribution of *An. oswaldoi* A now includes the Brazilian States of Acre, Amazonas, Mato Grosso, Pará and Rondônia, and the Colombian Department of Amazonas (Figure 4). Vector incrimination studies in Acre, Brazil determined *An. oswaldoi* as a local vector of malaria [53-56]. As the only confirmed species of the Oswaldoi Complex present in Acre, Brazil, it seems likely that *An. oswaldoi* A plays a role in malaria transmission in this state, and most probably throughout its distribution.

**Table 4 Retrospective identification of published COI sequences of *An. oswaldoi* s.l. (DQ784827-DQ784851 [16]) and *An. konderi* s.l. (JF437965-JF437974 [23]) in relation to species determined herein, through sequence homology**

Locality (Brazil)	GenBank accession	GenBank identification	osw s.s.	osw A	sp. nr. kon	kon of Sallum
Acre, Sena Madureira	DQ784827	<i>An. oswaldoi</i>			98%	
	DQ784828	<i>An. oswaldoi</i>			98%	
	DQ784829	<i>An. oswaldoi</i>			99%	
	DQ784830	<i>An. oswaldoi</i>			98%	
	DQ784831	<i>An. oswaldoi</i>			98%	
	DQ784832	<i>An. oswaldoi</i>	99%			
	DQ784833	<i>An. oswaldoi</i>	99%			
	DQ784834	<i>An. oswaldoi</i>	99%			
Amazonas, Coari	DQ784835	<i>An. oswaldoi</i>	99%			
	DQ784836	<i>An. oswaldoi</i>	99%			
	DQ784837	<i>An. oswaldoi</i>	99%			
Rondônia, São Miguel	DQ784838	<i>An. oswaldoi</i>	99%			
	DQ784839	<i>An. oswaldoi</i>				99%
	DQ784840	<i>An. oswaldoi</i>				99%
Rondônia, São Miguel	DQ784841	<i>An. oswaldoi</i>				99%
	DQ784842	<i>An. oswaldoi</i>				99%
	DQ784843	<i>An. oswaldoi</i>			100%	
	DQ784844	<i>An. oswaldoi</i>			99%	
	DQ784845	<i>An. oswaldoi</i>				99%
	DQ784846	<i>An. oswaldoi</i>				99%
	DQ784847	<i>An. oswaldoi</i>				99%
	DQ784848	<i>An. oswaldoi</i>				99%
Pará, Moju	DQ784849	<i>An. oswaldoi</i>		99%		
	DQ784850	<i>An. oswaldoi</i>		98%		
	DQ784851	<i>An. oswaldoi</i>		99%		
Acre, Acrelândia	JF437965	<i>An. konderi</i>			100%	
Amapá, Macapá	JF437966	<i>An. konderi</i>				100%
	JF437967	<i>An. konderi</i>				100%
	JF437968	<i>An. konderi</i>				99%
Rondônia, Monte Negro	JF437969	<i>An. konderi</i>	99%			
Paraná, Porto Natal	JF437970	<i>An. konderi</i>	99%			
	JF437971	<i>An. konderi</i>	99%			
	JF437972	<i>An. konderi</i>	99%			
	JF437973	<i>An. konderi</i>	99%			
Paraná, Santa Helena	JF437974	<i>An. konderi</i>	99%			

[osw s.s.: *An. oswaldoi* s.s.; osw A: *An. oswaldoi* A; sp. nr. kon: *An. sp. nr. konderi*; kon: *An. konderi* of Sallum from Macapá, Amapá, Brazil].

#### ***Anopheles* sp. nr. *konderi* (designated herein)**

ITS2 haplotype HXIII was shared by a number of individuals from Madre de Dios and Loreto provinces in Peru, Orellana province in Ecuador, and in the Department of Caquetá, southern Colombia (Figure 4). These sequences showed high similarity (98-100%) to specimens identified as *An. konderi* s.l. from Amapá, Brazil [GenBank: JF437934-36, JF437926] [23], differing by only two

transitions (G/A, position 361 and T/C, position 449) and a two base indels (at positions 264–265). We recognize that such intragenetic variation in ITS2 is not uncommon, having also been detected in *An. oswaldoi* A, *An. oswaldoi* B and other *Nyssorhynchus* species [9,57-59]. However, COI sequences from the same individuals analysed by Bayesian, MYC model and TCS network analyses confirm an independent sister cluster to

the *An. konderi* of Sallum from Macapá, Amapá, Brazil, also included in our analysis. Both these taxa are unique with respect to a third *An. konderi* s.l. from Acre reported by Sallum et al. [9] (GenBank: EU636801).

The type locality of *An. konderi* is Solimões River, Coari, Amazonas, Brazil. However, in their recent redescription of *An. konderi*, Flores-Mendoza et al. [21] were unable to access material from this locality, using several other localities in Brazil instead. Some samples used in our molecular analysis were the same as those examined by Flores-Mendoza et al. [21], but as the identity of *An. konderi* is uncertain, we have designated this cluster as *An. sp. nr. konderi*.

The trimmed 404 bp *An. oswaldoi* "Group 2" *COI* sequences of Scarpassa and Conn [16] (GenBank: DQ784827-31 and DQ784843-44) are 98-99% similar to our *An. sp. nr. konderi* and 100% identical to GenBank: JF437965 [16] (Table 4). Correlation of our data with published works confirms the presence of *An. sp. nr. konderi* in three localities in Brazil (Acrelândia and Sena Madureira (Acre) and São Miguel (Rondônia)), and our data suggest that *An. sp. nr. konderi* may be allopatric in Madre de Dios and Loreto provinces in Peru, where *An. oswaldoi* s.l. has been confirmed as an efficient malaria vector [60-62].

The taxonomic status of *An. konderi* s.l. clearly needs to be reassessed. This and other studies [9,16,23] have provided evidence for at least three species in *An. konderi* s.l. Concerted efforts are needed to acquire material from the type locality in order to establish the identity of *An. konderi* s.s. before the systematics of closely related taxa can be properly understood. Given the data presented here, it is likely that the species discussed here will fall into a more comprehensive Oswaldoi-Konderi Complex in future.

## Conclusions

Through novel data and retrospective comparison of *COI* and *ITS2* DNA data, evidence is shown to support three species within of *An. oswaldoi* s.l. (*An. oswaldoi* s.s., *An. oswaldoi* A and *An. oswaldoi* B), and at least three species in the closely related *An. konderi* s.l. (*An. sp. nr. konderi*, *An. konderi* of Sallum herein, and in Sallum et al. [9]). Determining the specific status and distribution of component members has allowed the circumstantial incrimination of *An. oswaldoi* A, *An. oswaldoi* B and *An. sp. nr. konderi* as malaria vectors. *Anopheles oswaldoi* s.s. has never been implicated in malaria transmission. Morphological studies of the species listed in the manuscript are now pertinent, to find reliable diagnostic characters, and to follow with the formal description and naming of the new species determined.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

FRL and YML conceived the ideas; YML, DJP, RCW and MLQ obtained funding; FRL, MLQ, MAMS, CFM, DC, JA, JAO and RCW undertook fieldwork and/or donated samples; FRL, YML and MH carried out the molecular laboratory work; FRL, YML and RCW carried the data analysis and interpretation; FRL wrote the draft manuscript; YML, RCW, MAMS, DJP, MLQ, IDV, CFM, and JA revised the draft manuscript; FRL, RCW and YML carried out the final revision and submitted the manuscript. All authors have read and approved the final version of the manuscript.

## Acknowledgements

This study formed part of the PhD study of FRL conducted at the Natural History Museum, London, and awarded from Canterbury Christ Church University, Canterbury, Kent, U.K. This investigation received financial support from the UNICEF/UNDP/World Bank/WHO Special Programme for Research and Training in Tropical Diseases (TDR) (grant A50252 to YML), Canterbury Christ Church University (studentship to FRL). Additional funding was obtained through the Friends of the Natural History Museum, London to further the activities of the Mosquito Barcoding Initiative (to YML); the Consortium for the Barcode of Life (CBOL) (to YML and RCW) and the Sloane Foundation (to YML and RCW); the National Institute of Health (NIH), USA (grant 2R01AI054139 to Jan E. Conn) and COLCIENCIAS (grant 110134319196 to MLQ). We thank Dr. A. Papadopoulos for help with the MYC analysis and Dr. S. Mahamdallie for helpful suggestions and discussions during preparation of the manuscript.

This manuscript was prepared in part whilst YML held a National Research Council Senior Research Associateship Award at the Walter Reed Army Institute of Research. This research was performed in part under a Memorandum of Understanding between the Walter Reed Army Institute of Research and the Smithsonian Institution, with institutional support provided by both organizations. The material to be published reflects the views of the authors and should not be construed to represent those of the Department of the Army or the Department of Defense.

## Author details

<sup>1</sup>Department of Entomology, National Museum of Natural History, Smithsonian Institution, Museum Support Center, Suitland, MD 20746, USA. <sup>2</sup>Programa de Estudio y Control de Enfermedades Tropicales (PECET), Facultad de Medicina, Universidad de Antioquia, Medellín, Colombia. <sup>3</sup>Walter Reed Biosystematics Unit, Smithsonian Institution, Museum Support Center, Suitland, MD 20746, USA. <sup>4</sup>Department of Geographical and Life Sciences, Canterbury Christ Church University, Kent, UK. <sup>5</sup>Facultad de Medicina, Universidad Nacional de Colombia, Bogotá, Colombia. <sup>6</sup>Departamento de Epidemiología, Faculdade de Saúde Pública, Universidade de São Paulo, São Paulo, SP, Brazil. <sup>7</sup>Naval Medical Research Unit (NAMRU-6), Lima, Peru. <sup>8</sup>Department of Life Science, Faculty of Science and Technology, University of the West Indies, St. Augustine Campus, West Indies, Trinidad and Tobago. <sup>9</sup>Servicio Nacional de Control de Enfermedades Transmitidas por Vectores Artrópodos, Ministerio Salud Pública, Guayaquil, Ecuador. <sup>10</sup>Walter Reed Army Institute of Research, 503 Robert Grant Avenue, Silver Spring, MD 20910, USA.

Received: 19 November 2012 Accepted: 10 September 2013

Published: 12 November 2013

## References

1. Harbach RE: The classification of genus *Anopheles* (Diptera: Culicidae): a working hypothesis of phylogenetic relationships. *Bull Entomol Res* 2004, **94**(6):537-553.
2. Bourke BP, Foster PG, Berge ES, Calado DC, Sallum MA: Phylogenetic relationships among species of *Anopheles* (*Nyssorhynchus*) (Diptera, Culicidae) based on nuclear and mitochondrial gene sequences. *Acta Trop* 2010, **114**(2):88-96.
3. Conn J: A genetic study of the malaria vector *Anopheles nuneztovari* from western Venezuela. *J Am Mosq Control Assoc* 1990, **6**(3):400-405.
4. Marrelli MT, Sallum MA, Marinotti O: The second internal transcribed spacer of nuclear ribosomal DNA as a tool for Latin American anopheline taxonomy: a critical review. *Mem Inst Oswaldo Cruz* 2006, **101**(8):817-832.

5. Motoki MT, Wilkerson RC, Sallum MA: **The *Anopheles albitarsis* complex with the recognition of *Anopheles oryzalimnetes* Wilkerson and Motoki, n. sp. and *Anopheles janconnae* Wilkerson and Sallum, n. sp. (Diptera: Culicidae).** *Mem Inst Oswaldo Cruz* 2009, **104**(6):823–850.
6. Rosa-Freitas MG, Lourenço-de-Oliveira R, de Carvalho-Pinto CJ, Flores-Mendoza C, Silva-do-Nascimento TF: **Anopheline species complexes in Brazil: current knowledge of those related to malaria transmission.** *Mem Inst Oswaldo Cruz* 1998, **93**(5):651–655.
7. Silva-do-Nascimento TF, Lourenço-de-Oliveira R: **Diverse population dynamics of three *Anopheles* species belonging to the triannulatus complex (Diptera: Culicidae).** *Mem Inst Oswaldo Cruz* 2007, **102**(8):975–982.
8. Ruiz-Lopez F, Wilkerson RC, Conn JE, McKeon SN, Levin DM, Quinones ML, Povoá MM, Linton Y-M: **DNA barcoding reveals both known and novel taxa in the Albitarsis group (*Anopheles*: *Nyssorhynchus*) of neotropical malaria vectors.** *Parasites and Vectors* 2012, **5**:44.
9. Sallum MA, Marrelli MT, Nagaki SS, Laporta GZ, Dos Santos CL: **Insight into *Anopheles (Nyssorhynchus)* (Diptera: Culicidae) species from Brazil.** *J Med Entomol* 2008, **45**(6):970–981.
10. Sierra DM, Velez ID, Linton Y-M: **Malaria vector *Anopheles (Nyssorhynchus) nuneztovari* comprises one genetic species in Colombia based on homogeneity of nuclear ITS2 rDNA.** *J Med Entomol* 2004, **41**(3):302–307.
11. Faran ME: **Mosquito studies (Diptera: Culicidae) XXXIV: a revision of the albitarsis section of the subgenus *Nyssorhynchus* of *Anopheles*.** *Contr Am Entomol Inst* 1980, **15**:1–215.
12. Motoki MT, Linton Y-M, Ruiz F, Flores-Mendoza C, Sallum MAM: **Redescription of *Anopheles oswaldoi* (Peryassú, 1922) (Diptera: Culicidae), with formal lectotype designation.** *Zootaxa* 2007, **1588**:31–51.
13. Quiñones ML, Ruiz F, Calle DA, Harbach RE, Erazo HF, Linton Y-M: **Incrimination of *Anopheles (Nyssorhynchus) rangeli* and *An. (Nys.) oswaldoi* as natural vectors of *Plasmodium vivax* in Southern Colombia.** *Mem Inst Oswaldo Cruz* 2006, **101**(6):617–623.
14. Marrelli MT, Malafronte RS, Flores-Mendoza C, Lourenço-de-Oliveira R, Kloetzel JK, Marinotti O: **Sequence analysis of the second internal transcribed spacer of ribosomal DNA in *Anopheles oswaldoi* (Diptera: Culicidae).** *J Med Entomol* 1999, **36**(6):679–684.
15. Marrelli MT, Floeter-Winter LM, Malafronte RS, Tadei WP, Lourenço-de-Oliveira R, Flores-Mendoza C, Marinotti O: **Amazonian malaria vector anopheline relationships interpreted from ITS2 rDNA sequences.** *Med Vet Entomol* 2005, **19**(2):208–218.
16. Scarpassa VM, Conn JE: **Molecular differentiation in natural populations of *Anopheles oswaldoi sensu lato* (Diptera: Culicidae) from the Brazilian Amazon, using sequences of the *COI* gene from mitochondrial DNA.** *Genet Mol Res* 2006, **5**(3):493–502.
17. Ruiz F, Quiñones ML, Erazo HF, Calle DA, Alzate JF, Linton Y-M: **Molecular differentiation of *Anopheles (Nyssorhynchus) benarrochi* and *An. (N.) oswaldoi* from southern Colombia.** *Mem Inst Oswaldo Cruz* 2005, **100**(2):155–160.
18. Quiñones ML, Harbach RE, Calle DA, Ruiz F, Erazo HF, Linton Y-M: **ariante morfológica de adultas hembras de *Anopheles benarrochi* (Diptera: Culicidae) en Putumayo, Colombia.** *Biomedica* 2001, **21**:351–359.
19. Galvão ALA, Damasceno RG: ***Anopheles (Nyssorhynchus) konderi* nova espécie de *Anopheles* do vale do Amazonas e considerações sobre as espécies do complexo *Tarsimaculatus* (Diptera, Culicidae).** *Folia Clin Biol* 1942, **14**(5–6):115–135.
20. Lane J: **Neotropical culicidae.** *Universidade de São Paulo* 1953, **1**:548pp.
21. Flores-Mendoza C, Peyton EL, Wilkerson RC, Lourenço-de-Oliveira R: ***Anopheles (Nyssorhynchus) konderi* Galvão and Damasceno: neotype designation and resurrection from synonymy with *Anopheles (Nyssorhynchus) oswaldoi* (Peryassú) (Diptera: Culicidae).** *Proc Entomol Soc Wash* 2004, **106**:118–132.
22. Scarpassa VM: **Isozyme similarity in *Anopheles oswaldoi sensu lato* (Diptera: Culicidae) from the Amazon Region, Brazil.** *J Med Entomol* 2005, **42**(3):319–326.
23. Motoki MT, Bourke BP, Berço ES, Da Silva AM, Sallum MAM: **Systematic notes of *Anopheles konderi* and its first record in Paraná State, Brazil.** *J Am Mosq Control Assoc* 2011, **27**(3):191–200.
24. Motoki MT, Dos Santos CLS, Sallum MAM: **Intraspecific variation on the aedeagus of *Anopheles oswaldoi* (Peryassú) (Diptera: Culicidae).** *Neotrop Entomol* 2009, **38**(1):890–894.
25. Pinault LL, Hunter FF: **New highland distribution records of multiple *Anopheles* species in the Ecuadorian Andes.** *Malaria J* 2011, **10**:236.
26. Foley D: **A spreadsheet mapping approach for error checking and sharing collection point data.** *Biodiv Inform* 2011, **7**:137–142.
27. Collins FH, Paskewitz SM: **A review of the use of ribosomal DNA (rDNA) to differentiate among cryptic *Anopheles* species.** *Insect Mol Biol* 1996, **5**(1):1–9.
28. Linton Y-M, Harbach RE, Chang MS, Anthony TG, Matusop A: **Morphological and molecular identity of *Anopheles (Cellia) sundaicus* (Diptera: Culicidae), the nominotypical member of a malaria vector species complex in Southeast Asia.** *Syst Ent* 2001, **26**:357–366.
29. Hebert PDN, Cywinska A, Ball SL, deWaard JR: **Biological identifications through DNA barcodes.** *Proc Biol Sci* 2003, **270**(1512):313–321.
30. Folmer O, Black M, Hoeh W, Lutz R, Vrijenhoek R: **DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates.** *Mol Mar Biol Biotechnol* 1994, **3**(5):294–299.
31. Ruiz F, Linton Y-M, Ponsonby DJ, Conn JE, Herrera M, Quiñones ML, Velez ID, Wilkerson RC: **Molecular comparison of topotypic specimens confirms *Anopheles (Nyssorhynchus) dunhami* Causey (Diptera: Culicidae) in the Colombian Amazon.** *Mem Inst Oswaldo Cruz* 2010, **105**(7):899–903.
32. Maddison DR, Maddison WR: **MacClade: Analysis of Phylogeny and Character Evolution.** Sunderland: Sinauer Associates; 2003.
33. Clement M, Posada D, Crandall KA: **TCS: a computer program to estimate gene genealogies.** *Mol Ecol* 2000, **9**(10):1657–1659.
34. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S: **MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods.** *Mol Biol Evol* 2011, **28**(10):2731–2739.
35. Kimura M: **A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences.** *J Mol Evol* 1980, **16**(2):111–120.
36. Nylander J: **MrModeltest v2.** Program distributed by the author: Evolutionary Biology Center, Uppsala University; 2004.
37. Ronquist F, Huelsenbeck JP: **MrBayes 3: Bayesian phylogenetic inference under mixed models.** *Bioinformatics* 2003, **19**(12):1572–1574.
38. Swofford DL: **PAUP\*: Phylogenetic Analysis Using Parsimony (\*and Other Methods): Version 4.** Sunderland, Massachusetts: Sinauer Associates, Inc.; 2002.
39. Felsenstein J: **Confidence limits on phylogenies: an approach using the bootstrap.** *Evolution* 1985, **39**:783–791.
40. Fontaneto D, Herniou EA, Boschetti C, Caprioli M, Melone G, Ricci C, Barraclough TG: **Independently evolving species in asexual bdelloid rotifers.** *PLoS Biol* 2007, **5**(4):e87.
41. Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, Kamoun S, Sumlin WD, Vogler AP: **Sequence-based species delimitation for the DNA taxonomy of undescribed insects.** *Syst Biol* 2006, **55**(4):595–609.
42. Yule GU: **A mathematical theory of evolution, based on the conclusions of Dr. J.C. Willis, FRS.** *Phil Trans Roy Soc London Series B* 1925, **213**:21–87.
43. Sanderson MJ: **r8s: inferring absolute rates of molecular evolution and divergence times in the absence of a molecular clock.** *Bioinformatics* 2003, **19**(2):301–302.
44. Rambaut A: **FigTree. v1.3.1.** ; 2009 (Available for download from <http://tree.bio.ed.ac.uk/software/figtree>).
45. Sites JW, Marshall JC: **Delimiting species: a Renaissance issue in systematic biology.** *Trends Ecol Evol* 2003, **18**:462–470.
46. de Oliveira-Ferreira J, Lourenço-de-Oliveira R, Teva A, Deane LM, Daniel-Ribeiro CT: **Natural malaria infections in anophelines in Rondônia State, Brazilian Amazon.** *Am J Trop Med Hyg* 1990, **43**(1):6–10.
47. Klein TA, Lima JB, Tada MS: **Comparative susceptibility of anopheline mosquitoes to *Plasmodium falciparum* in Rondônia, Brazil.** *Am J Trop Med Hyg* 1991, **44**(6):598–603.
48. Klein TA, Lima JB, Tada MS, Miller R: **Comparative susceptibility of anopheline mosquitoes in Rondônia, Brazil to infection by *Plasmodium vivax*.** *Am J Trop Med Hyg* 1991, **45**(4):463–470.
49. Povoá MM, Wirtz RA, Lacerda RNL, Miles MA, Warhurst D: **Malaria vectors in the municipality of Serra do Navio, State of Amapá Amazon Region, Brazil.** *Mem Inst Oswaldo Cruz* 2001, **96**(2):179–184.
50. Rubio-Palis Y, Curtis CF: **Biting and resting behaviour of anophelines in western Venezuela and implications for control of malaria transmission.** *Med Vet Entomol* 1992, **6**(4):325–334.
51. Rozeboom LE: **Subspecific variations among neotropical *Anopheles* mosquitoes, and their importance in the transmission of malaria.** *American J Trop Med Hy* 1942, **22**(3):235–255.
52. Curry DP: **Some observations on the *Nyssorhynchus* group of the *Anopheles* (Culicidae) of Panama.** *Am J Epidemiol* 1932, **15**(2):566–572.
53. Branquinho MS, Lagos CBT, Rocha RM, Natal D, Barata JM, Cochrane AH, Nardin E, Nussenzeig RS, Kloetzel JK: **Anophelines in the state of Acre,**

- Brazil, infected with *Plasmodium falciparum*, *P. vivax*, the variant *P. vivax* VK247 and *P. malariae*. *Trans R Soc Trop Med Hyg* 1993, **87**(4):391–394.
54. Branquinho MS, Araujo MS, Natal D, Marrelli MT, Rocha RM, Taveira FAL, Kloetzel JK: *Anopheles oswaldoi* a potential malaria vector in Acre, Brazil. *Trans R Soc Trop Med Hyg* 1996, **90**(3):233.
  55. Marrelli MT, Branquinho MS, Hoffmann EHE, Taipe-Lagos CB, Natal D, Kloetzel JK: Correlation between positive serology for *Plasmodium vivax*-like/*Plasmodium simiovale* malaria parasites in the human and anopheline populations in the State of Acre, Brazil. *Trans R Soc Trop Med Hyg* 1998, **92**(2):149–151.
  56. Marrelli MT, Honorio NA, Flores-Mendoza C, Lourenço-de-Oliveira R, Marinotti O, Kloetzel JK: Comparative susceptibility of two members of the *Anopheles oswaldoi* complex, *An. oswaldoi* and *An. konderi*, to infection by *Plasmodium vivax*. *Trans R Soc Trop Med Hyg* 1999, **93**(4):381–384.
  57. Fairley TL, Kilpatrick CW, Conn JE: Intragenomic heterogeneity of internal transcribed spacer rDNA in neotropical malaria vector *Anopheles aquasalis* (Diptera: Culicidae). *J Med Entomol* 2005, **42**(5):795–800.
  58. Li C, Wilkerson RC: Intragenomic rDNA ITS2 variation in the neotropical *Anopheles (Nyssorhynchus) albitarsis* complex (Diptera: Culicidae). *J Hered* 2007, **98**(1):51–59.
  59. Onyabe DY, Conn JE: Intragenomic heterogeneity of a ribosomal DNA spacer (ITS2) varies regionally in the neotropical malaria vector *Anopheles nuneztovari* (Diptera: Culicidae). *Insect Mol Biol* 1999, **8**(4):435–442.
  60. Calderón G, Fernández R, Valle J: Especies de la fauna anofelina, su distribución y algunas consideraciones sobre su abundancia e infectividad en el Perú. *Revista Peruana de Epidemiología* 1995, **8**(2):5–23.
  61. Hayes J, Calderon G, Falcon R, Zambrano V: Newly incriminated anopheline vectors of human malaria parasites in Junin Department, Peru. *J Am Mosq Control Assoc* 1987, **3**(3):418–422.
  62. Guarda JA, Asayag CR, Witzig R: Malaria reemergence in the Peruvian Amazon region. *Emerg Infect Dis* 1999, **5**(2):209–215.

doi:10.1186/1756-3305-6-324

Cite this article as: Ruiz-Lopez *et al.*: Systematics of the Oswaldoi Complex (*Anopheles*, *Nyssorhynchus*) in South America. *Parasites & Vectors* 2013 **6**:324.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

