

ERRATUM

Open Access



Erratum to: Physician reported incidence of early and late Lyme borreliosis

Agnetha Hofhuis*, Margriet Harms, Sita Bennema, Cees C. van den Wijngaard and Wilfrid van Pelt

After the publication of this work [1], we noticed an error in the percentages of “Fig. 1 Lyme borreliosis manifestations as a proportion of all Lyme-related diagnoses, by type of physician”. The percentages of the first horizontal bar “General Practitioner (adjusted)” should be: 90.9 % erythema migrans (instead of 92 %), 5.3 % disseminated Lyme borreliosis, and 3.8 % persisting symptoms attributed to Lyme borreliosis (instead of 3 %).

Consequently, the corresponding sentence in the results section should be: “After adjusting, erythema migrans represented 90.9 % of the clinical diagnoses for Lyme borreliosis reported by the GP, while 5.3 % of the diagnoses concerned disseminated Lyme borreliosis, and 3.8 % concerned persisting symptoms attributed to Lyme borreliosis (see Fig. 1)”.

And the corresponding sentence in the discussion section should be: “Considering all GP-reported Lyme borreliosis, 5.3 % of the diagnoses concerned disseminated Lyme borreliosis, and 3.8 % concerned persisting symptoms attributed to Lyme borreliosis (see Fig. 1).”

Received: 6 July 2015 Accepted: 6 July 2015

Published online: 17 July 2015

Reference

1. Hofhuis A, Harms M, Bennema S, van den Wijngaard CC, van Pelt W. Physician reported incidence of early and late Lyme borreliosis. *Parasit Vectors*. 2015;8:161.

* Correspondence: Agnetha.Hofhuis@rivm.nl
Epidemiology and Surveillance Unit, Centre for Infectious Disease Control Netherlands, National Institute for Public Health and the Environment (RIVM), Bilthoven, The Netherlands

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit






Fig. 1 Lyme borreliosis manifestations as a proportion of all Lyme-related diagnoses, by type of physician